arXiv:1110.1874v3 [math.DG] 28 Nov 2011

MAXIMUM RANK OF A LEGENDRIAN WEB

JOE S. WANG

ABSTRACT. We propose the Legendrian web in a contact three manifold for a second order generalization of the planar web. An Abelian relation for a Legendrian web is analogously defined as an additive relation among the first integrals of foliations. For a class of Legendrian d-webs defined by simple second order ODE's, we give an algebraic construction of $\rho_d = \frac{(d-1)(d-2)(2d+3)}{6}$ linearly independent Abelian relations. We then employ the method of local differential analysis and show that ρ_d is the maximum rank of a Legendrian d-web.

The structure equations for the Legendrian 3-webs of maximum rank three are determined, and their explicit local normal forms are given. For an application, we give an alternative characterization of a Darboux super-integrable metric as a two dimensional Riemannian metric g_+ that admits a mate metric g_- such that a Legendrian 3-web naturally associated with the geodesic foliations of the pair $g_\pm$ has maximum rank.

1. INTRODUCTION

A *d-web of codimension ν* in a manifold is a set of d foliations by submanifolds of codimension ν whose d tangent spaces are in general position at each point, [10]. Owing to the basic nature of definition, the web structure can be found in diverse areas of differential geometry, [2][13][11][18][27][1][30].

As pointed out by Chern in the survey [13], Lie's theorem on the surfaces of double translation can be restated in web geometry terms that a planar 4-web of curves of maximum rank three(see below) is algebraic, which is also related to a particular case of the converse of Abel's theorem. For an example in projective geometry, the classical Chasles' theorem on the intersecting cubics is linked with the hexagonality of algebraic planar 3-webs. According to [1], a three dimensional semi-simple Frobenius manifold carries a family of characteristic hexagonal planar 3-webs. Planar 3-webs also naturally arise as the asymptotic webs of Legendrian surfaces in the projective space $\mathbb{P}^5$, [30]. Although in a possibly different context, the author's own interests in web geometry originated from the Gelfand & Zakharevich's work on the Veronese webs attached to bi-Hamiltonian systems, [18]. Much of the developments of web geometry up until 1930's are recorded in the Blaschke & Bol's book "Geometrie der Gewebe", [2]. For the more recent literature on the subject, we refer to [17][27] and the references therein.

Let $\mathcal{W}$ be a d-web of codimension ν. Since a single regular foliation (of any fixed codimension) in a finite dimensional manifold has no local invariants, the local geometry of $\mathcal{W}$ lies in the relative position among the d foliations. A pertinent notion in this aspect is the *Abelian k-relation(equation)*, $1 \leq k \leq \nu$, which represents the additive relation among the closed basic k-forms with respect to each foliations.[1] The set of Abelian k-relations of $\mathcal{W}$ naturally form a vector space, and the *k-rank* of $\mathcal{W}$ denoted by $r^k(\mathcal{W})$ is the dimension of the vector space of its Abelian k-relations. The sequence of numbers $r^k(\mathcal{W})$, $k = 1, 2, \ldots \nu$, is the basic invariant of a web.

The k-rank of a d-web of codimension ν is not arbitrary. There generally exists an optimal bound determined in terms of k, d, ν, and the dimension of the ambient space, [11][19]. It is a classical result of Bol that the maximum rank(1-rank) for a planar d-web is $\frac{(d-1)(d-2)}{2}$, which is the well known genus bound for a degree d plane algebraic curve, [5]. Chern in his thesis generalized this and showed that the maximum rank for a d-web of codimension 1 is equal to the Castelnuovo bound for a degree d algebraic curve, [12].

2000 *Mathematics Subject Classification.* 53A60.

Key words and phrases. contact three manifold, Legendrian web, Abelian relation.

[1] An Abelian 1-relation represents an additive relation among the (differentiated) first integrals of the foliations.




Figure 1.1. Projective duality

The idea behind these results is the projective duality, Fig. 1.1, and the corresponding Abel's theorem, [10].[2] A degree d curve $\Sigma \subset (\mathbb{P}^n)^*$ induces by the standard dual construction an associated d-web of hyperplanes $\mathcal{W}_\Sigma$ in $\mathbb{P}^n$. By Abel's theorem, the trace of a holomorphic 1-form on Σ gives an Abelian relation for $\mathcal{W}_\Sigma$. The dimension of the subspace of such algebraic Abelian relations for $\mathcal{W}_\Sigma$ is then bounded by the Castelnuovo bound for an algebraic curve.

With this background geometry, the results of Bol, and Chern admit the following interpretation. The vector space of Abelian relations (for a web of codimension 1) is by definition the space of solutions to a linear differential system for the sections of a direct sum bundle canonically associated with a web. The differential system is generally over-determined, see Section 2. It is not difficult to imagine that such a uniform rank bound, if exists, is determined entirely by the symbol of the linear differential system. The results of Bol, and Chern indicate that even when one considers the general webs of codimension 1, and not just the kind of algebraic webs $\mathcal{W}_\Sigma$ described above, the relevant symbol of the linear differential system for Abelian relations is isomorphic in an appropriate sense to that of the algebraic webs. The canonical linear differential system for Abelian relations does not get any more over-determined in symbol-wise when one extends the category from algebraic webs to general webs.

We shall consider in this paper a generalization of web geometry to the geometric situation where the foliations are subject to a first order constraint. Let X be a three dimensional manifold. A *contact structure* on X is a 2-plane field $\mathcal{D} \subset TX$ locally defined by $\mathcal{D} = \langle\, \theta \,\rangle^\perp$ for a 1-form θ that satisfies the non-degeneracy condition $d\theta \wedge \theta \neq 0$. Given a contact structure, a *Legendrian curve* is an immersed D-horizontal curve. A *Legendrian d-web* is a set of d pairwise transversal foliations by Legendrian curves. The Pfaff-Darboux local normal form theorem for contact structures shows that a Legendrian d-web is locally defined by a set of d second order ODE's.

Recall that a planar d-web is a set of d transversal foliations by curves on a two dimensional manifold, which is locally defined by a set of d first order ODE's. The purpose of this paper is to propose the Legendrian web in a contact three manifold for a second order generalization of the planar web. An Abelian relation, which represents an additive functional relation among the first integrals of foliations, is analogously defined for the Legendrian web. The rank of a Legendrian web is the dimension of the vector space of Abelian relations.

In analogy with the discussion above, the complex algebraic model for our investigation is the duality associated with the symplectic group $\mathrm{Sp}_2\mathbb{C}$, Fig. 1.2. Here the 3-quadric $\mathbb{Q}^3$ is the space of Legendrian lines in $\mathbb{P}^3$, and dually $\mathbb{P}^3$ is the space of null lines in $\mathbb{Q}^3$ with respect to the $\mathrm{Sp}_2\mathbb{C}$ invariant contact, and conformal structures respectively. Z is the incidence space, [8]. Consider a degree $2d$ analytic surface $\Sigma \subset \mathbb{Q}^3 \subset \mathbb{P}^4$. Then generically Σ intersects a null line at d distinct points, and it induces by duality an associated d-web $\mathcal{W}_\Sigma$ of Legendrian lines on an open subset of $\mathbb{P}^3$. This suggests to consider a Legendrian d-web as *a generalized degree $2d$ surface in $\mathbb{Q}^3 \subset \mathbb{P}^4$*.

Let Ω be a closed meromorphic 1-form on Σ. In consideration of the Abel's theorem, one may insist that Ω is *holomorphic* when the trace $(\pi_1)_* \circ (\pi_2)^*\Omega$ vanishes.[3] By definition of the trace, such a closed holomorphic 1-form gives rise to an Abelian relation for $\mathcal{W}_\Sigma$. An Abelian relation for a Legendrian d-web can therefore be considered as *a generalized closed holomorphic 1-form on a degree $2d$ surface in $\mathbb{Q}^3 \subset \mathbb{P}^4$*.

One of the initial motivations for the present work came from the observation that the bound for the rank of a Legendrian d-web would imply the Castelnuovo type bound for a degree $2d$ surface in $\mathbb{Q}^3$; determining the rank

[2] Many of the arguments and results in web geometry hold both in the real smooth, and the complex analytic categories. Let us argue here in the complex analytic category.

[3] The relevant technical details are being ignored here. Since we are attempting to define holomorphic objects on generally singular surfaces, there are a few different choices, [20]. Note also that a holomorphic 1-form on a smooth surface is necessarily closed.




Figure 1.2. Projective duality associated with $\mathrm{Sp}_2\mathbb{C}$

bound by a direct local analysis of the linear differential system for Abelian relations, one may derive conversely a global Castelnuovo bound for the analytic surfaces in $\mathbb{Q}^3$.

Main results.

1. We give an algebraic construction of $\frac{(d-1)(d-2)(2d+3)}{6}$ linearly independent Abelian relations for a class of simple Legendrian d-webs, Section 3.

2. The maximum rank of a Legendrian d-web is

$$\rho_d = \frac{(d-1)(d-2)(2d+3)}{6}. \tag{1.1}$$

3. We give a local analytic characterization of the Legendrian 3-webs of maximum rank three,[4] and determine their local normal forms, Proposition 5.6, Theorem 5.9. We give an alternative characterization of a Darboux super-integrable metric in terms of a Legendrian 3-web associated with its geodesic flow, Theorem 5.18.

In addition to the works of Bol, and Chern cited above, Chern & Griffiths determined the bound for the ν-rank of the webs of codimension ν in a manifold of dimension νm, [11]. Hénaut further generalized Chern & Griffiths' bound to the bound for the k-rank for $1 \leq k \leq \nu$. We refer to [14][22][26][28] for the related works on the webs of maximum rank. **Main results** can be considered as a partial generalization of these works to the Legendrian web.

The rank bound in Main result 2 calls for the study of the corresponding extremal degree $2d$ surfaces in $\mathbb{Q}^3$ with ρ_d holomorphic 1-forms.

The paper is organized as follows. In Section 1.1, we review the basic results on the contact structure on a three dimensional manifold. In Section 2, the main subjects of the paper are introduced, Legendrian webs and Abelian relations. In Section 3, we give an algebraic construction of ρ_d linearly independent Abelian relations for a class of Legendrian d-webs, (3.1), Theorem 3.3. Section 4 contains the main technical part of the paper. We first determine the infinite prolongation of the structure equation for the first integrals of a Legendrian foliation, Lemma 4.12. The prolonged structure equation is applied to the defining additive equation of Abelian relation, and yields an infinite sequence of higher order compatibility equations, Lemma 4.17. Two different approaches to the proof of rank bound are proposed. An analysis of a part of the symbol of the compatibility equations shows that the proof of rank bound can be reduced to checking that a sequence of integer matrices are non-degenerate in a strict sense, Theorem 4.24. The rank bound Theorem 4.37 is obtained by examining the whole symbol of the compatibility equations with respect to the Abelian relations constructed in Section 3. In Section 5, we give an analytic characterization of the Legendrian 3-webs of maximum rank three, Proposition 5.6, Theorem 5.9. We apply this to the Legendrian 3-webs associated with a pair of Riemannian metrics, Section 5.2. We give an alternative characterization of a Darboux super-integrable metric in terms of a Legendrian web associated with its geodesic flow, Theorem 5.18. In Section 6, we give three remarks on the related problems. Appendix contains the `Maple` code to check our claim on the non-degeneracy of a sequence of universal integer coefficient matrices.

We assume the real smooth category. The analysis and results are valid within the complex analytic category with minor modifications. All of the arguments in this paper are local.

The over-determined PDE machinery are used throughout the paper without specific reference. For the standard reference, we cite [3][21].

[4] The maximum rank of a 3-web of curves in a three dimensional manifold is five, [2, Section 36, 37].

We shall frequently use the following property of a Vandermonde matrix; given a set of d distinct numbers q^a, $a = 1, 2, \ldots d$, the Vandermonde matrix $(V_a^b) = ((q^a)^{b-1})_{a,b=1}^d$ has full rank.

1.1. **Contact three manifold.** In this section, we record the Pfaff-Darboux local normal form theorem for contact structures on a three dimensional manifold. The Pfaff-Darboux theorem is a basic structure theorem which gives a local uniformization of contact structures. For our purpose, it provides an analytic representation of a Legendrian web in terms of a finite set of second order ODE's for one scalar function of one variable.

Let X be a three dimensional manifold. Let $TX \to X$ be the tangent bundle, and let $T^*X \to X$ be the cotangent bundle of X.

Definition 1.2. *A* contact structure *on a three dimensional manifold X is a 2-plane field $\mathcal{D} \subset TX$ such that it is locally defined by $\mathcal{D} = \langle\, \theta \,\rangle^\perp$ for a 1-form θ that satisfies the non-degeneracy condition*

$$d\theta \wedge \theta \neq 0. \tag{1.3}$$

The contact line bundle *$\mathcal{C} \subset T^*X$ is the dual line bundle $\mathcal{C} = \mathcal{D}^\perp$. A* contact three manifold *is a three dimensional manifold with a given contact structure. A* (local) contact transformation *of a contact three manifold is a (local) diffeomorphism which preserves the contact structure.*

The contact three manifold in the following example will be used repeatedly for the rest of the paper.

Example 1.4. *The standard model of a contact three manifold is the first jet space $J^1(\mathbb{R}, \mathbb{R})$ of real valued functions on $\mathbb{R}$. Let x be the coordinate of $\mathbb{R}$, and let y, p denote the real valued function y of x, and the first derivative $p = \frac{dy}{dx}$ respectively. (x, y, p) is the standard coordinate of $J^1(\mathbb{R}, \mathbb{R})$. The contact line bundle is generated by the canonical contact 1-form*

$$\theta = dy - p\, dx. \tag{1.5}$$

Consider the contact transformation $(x, y, p) \to (p, y - px, -x)$. The full group of contact transformations is strictly larger than the group of transformations of $J^1(\mathbb{R}, \mathbb{R})$ induced by the point transformations, $(x, y, p) \to (\hat{x}(x, y), \hat{y}(x, y), \frac{\hat{y}_x + p\hat{y}_y}{\hat{x}_x + p\hat{x}_y})$.

There exists a distinguished class of curves on a contact three manifold.

Definition 1.6. *Let X be a contact three manifold with the contact structure $\mathcal{D} \subset TX$. A* Legendrian curve *is an immersed $\mathcal{D}$-horizontal curve.*

From the non-degeneracy condition (1.3), it is clear that there do not exist immersed $\mathcal{D}$-horizontal surfaces.

Example 1.7 (*continued from Example 1.4*)**.** *By definition of the canonical contact 1-form* (1.5)*, the first jet graph $(x, y = f(x), p = f'(x)) \subset J^1(\mathbb{R}, \mathbb{R})$ of an arbitrary real valued function $f(x)$ is a Legendrian curve.*

The following Pfaff-Darboux local normal form theorem states that a Legendrian curve in a contact three manifold is locally isomorphic to Example 1.7.

Pfaff-Darboux local normal form *Let X be a contact three manifold. Let $\mathrm{x}_0 \in X$ be a reference point. There exists an adapted local coordinate (x, y, p) of X in a neighborhood of x_0 such that the contact line bundle is generated by*

$$\mathcal{C} = \langle\, dy - p\, dx \,\rangle. \tag{1.8}$$

Let $\gamma \hookrightarrow X$ be a Legendrian curve through x_0. Then the adapted local coordinate (x, y, p) can be chosen such that

$$\begin{aligned} (x, y, p) &= (0, 0, 0), \quad \textit{at} \quad \mathrm{x}_0, \\ (y, p) &= (0, 0), \quad \textit{along} \quad \gamma. \end{aligned}$$

Proof. The existence of an adapted coordinate $(x,\, y,\, p)$ for (1.8) is well known, [3, p38]. Given the reference point $\mathrm{x}_0 \in X$, let $(x,\, y,\, p)|_{\mathrm{x}_0} = (x_0,\, y_0,\, p_0)$. From

$$dy - pdx = d(y - p_0x - y_0 + p_0x_0) - (p - p_0)d(x - x_0),$$

one may take $(\hat{x},\, \hat{y},\, \hat{p}) = (x - x_0,\, y - p_0x - y_0 + p_0x_0,\, p - p_0)$ as the desired coordinate adapted at x_0.

Let $(x,\, y,\, p)$ be a coordinate system adapted at x_0 as above. Given a Legendrian curve γ through x_0, there are three cases (they are not mutually exclusive).

• case $dx|_\gamma \neq 0$ at x_0.

Let $\gamma^*y = f(x)$, $\gamma^*p = f'(x)$ locally in a neighborhood of x_0. From $dy - pdx = d(y - f(x)) - (p - f'(x))dx$, one may take $(\hat{x},\, \hat{y},\, \hat{p}) = (x,\, y - f(x),\, p - f'(x))$ as the desired coordinate adapted along γ.

• case $dy|_\gamma \neq 0$ at x_0.

One may first translate $(x,\, y,\, p) \to (x,\, y + x,\, p + 1)$ so that $(x,\, y,\, p)|_{\mathrm{x}_0} = (0,\, 0,\, 1)$. The contact 1-form $dy - pdx$ can then be written as $dy - pdx = -p(dx - \frac{1}{p}dy)$. By interchanging x and y, and up to a translation similar as above, this is reduced to the first case.

• case $dp|_\gamma \neq 0$ at x_0.

Under the contact transformation $(x,\, y,\, p) \to (p,\, y - px,\, -x)$ in Example 1.4, this is reduced to the first case. □

The Pfaff-Darboux local normal form theorem reflects the ampleness of the local group of contact transformations. We will see in the next section(Lie's theorem) that the local group of contact transformations is in fact large enough to normalize not just a single Legendrian curve but a foliation by Legendrian curves.

2. Legendrian web

In this section, we introduce the main subjects of this paper, Legendrian web and its Abelian relations. In Section 2.1, a Legendrian web is defined as a finite set of foliations by Legendrian curves. In Section 2.2, an Abelian relation for a Legendrian web is defined as an additive relation among the first integrals of the foliations.

We continue the analysis of Section 1.1.

2.1. **Legendrian web.**

Definition 2.1. *A* Legendrian d-web *on a contact three manifold is a set of d pairwise transversal foliations by Legendrian curves.*

Let X be a contact three manifold. Let $\mathcal{W}$ be a Legendrian d-web on X. Let $\mathcal{C} \subset T^*X$ be the contact line bundle, and let $\mathcal{I}^a \subset T^*X$, $a = 1,\, 2,\, ...\, d$, be the d dual rank 2 sub-bundles which define $\mathcal{W}$. From the transversality condition, $\mathcal{I}^a \cap \mathcal{I}^b = \mathcal{C}$ for $a \neq b$. By the Pfaff-Darboux local normal form theorem, after modifying the adapted coordinate $(x,\, y,\, p)$ if necessary as was done in the proof of the theorem in Section 1.1, there exist an adapted local coordinate $(x,\, y,\, p)$ of X and a set of d functions $q^a(x, y, p)$, $a = 1,\, 2,\, ...\, d$; $q^a \neq q^b$ for $a \neq b$, such that

$$\mathcal{I}^a = \langle\, dy - p\, dx,\, dp - q^a\, dx\,\rangle, \quad a = 1,\, 2,\, ...\, d.$$

The geometry of a Legendrian d-web $\mathcal{W}$ is thus locally equivalent to the geometry of a set of d second order ODE's

$$y'' = q^a(x,\, y,\, y'), \quad a = 1,\, 2,\, ...\, d, \tag{2.2}$$

up to contact transformation.

Example 2.3. *A theorem of Lie states that a second order ODE for one scalar function of one variable is locally equivalent to the equation $y'' = 0$ up to contact transformation,* [25]. *The corresponding rank 2 sub-bundle $\mathcal{I}$ is generated by $\mathcal{I} = \langle\, dy - p\, dx,\, dp\,\rangle$. Under the contact transformation $(x,\, y,\, p) \to (p,\, y - px,\, -x)$, one gets*

$$\mathcal{I} = \langle\, dy,\, dx\,\rangle.$$

Lie's theorem implies that a Legendrian 1-web is locally equivalent to the foliation by the fibers of the natural projection $J^1(\mathbb{R},\, \mathbb{R}) \to J^0(\mathbb{R},\, \mathbb{R}) \simeq \mathbb{R}^2$.

Since a contact transformation that preserves these fibers of the projection is locally induced from a point transformations of $\mathbb{R}^2$, it follows that the geometry of a Legendrian d-web is locally equivalent to the geometry of a set of $d-1$ second order ODE's up to point transformation. In this respect, Cartan's work [9] on second order ODE's can be considered as a study of Legendrian 2-webs. Considering the main theme of the paper, the Abelian relations for a Legendrian web, we shall restrict our attention to the case of Legendrian d-webs for $d \geq 3$, see the argument below Definition 2.7.

Let us introduce a geometrical construction for Legendrian webs.

Example 2.4. *Let $J^1(\mathbb{R},\, \mathbb{R}) \to J^0(\mathbb{R},\, \mathbb{R}) \simeq \mathbb{R}^2$ be the natural projection from the 1-jet space $J^1(\mathbb{R},\, \mathbb{R})$ equipped with the canonical contact structure. Let $(x,\, y,\, p)$ be the standard adapted coordinate of $J^1(\mathbb{R},\, \mathbb{R})$ so that the projection map is given by $(x,\, y,\, p) \to (x,\, y)$, Example 1.4. Consider a two parameter family of curves in $\mathbb{R}^2$, each of which is locally a graph of a function $y = y(x)$. Assuming this family of curves are sufficiently generic to second order, it is clear that their Legendrian lifts $(x,\, y(x),\, y'(x))$ to $J^1(\mathbb{R},\, \mathbb{R})$ define a Legendrian foliation on an open subset of $J^1(\mathbb{R},\, \mathbb{R})$.*

Example 2.5 (*continued*)**.** *Let $\{\, \mathrm{p}^1,\, \mathrm{p}^2,\, ...\, \mathrm{p}^d \,\} \subset \mathbb{R}^2$ be a set of d distinct points. Consider for each p^a the two parameter pencil of y as quadratic polynomial of x based at p^a. A set of d distinct points in $\mathbb{R}^2$ in this way give rise to a Legendrian d-web on an open subset of $J^1(\mathbb{R},\, \mathbb{R})$.*

Example 2.6 (*continued*)**.** *Let $\{\, \mathrm{p}^1,\, \mathrm{p}^2,\, \mathrm{p}^3 \,\} \subset \mathbb{R}^2$ be a set of three distinct points. In addition to the quadratic pencil 3-web based at these points in Example 2.5, one may consider the Legendrian foliation induced by the two parameter family of conics through the three points. A set of three distinct points in $\mathbb{R}^2$ in this way give rise to a Legendrian 4-web on an open subset of $J^1(\mathbb{R},\, \mathbb{R})$. It is a Legendrian analogue of Bol's exceptional 5-web,* [14].

2.2. **Abelian relation.** In planar web geometry, the projective duality suggests to consider a planar d-web as a generalized degree d plane curve. An Abelian relation for a planar d-web formally corresponds to a holomorphic 1-form through Abel's theorem, [10].

We follow this line of idea and give a definition for the Abelian relation of a Legendrian web.

Definition 2.7. *Let X be a contact three manifold. Let $\mathcal{W}$ be a Legendrian d-web on X. Let $\mathcal{I}^a \subset T^*X$, $a = 1,\, 2,\, ...\, d$, be the dual rank 2 sub-bundles which define $\mathcal{W}$. An* Abelian relation *of $\mathcal{W}$ is a d-tuple of 1-forms*[5] *$(\Omega^1,\, \Omega^2,\, ...\, \Omega^d)$, $\Omega^a \in C^\infty(\mathcal{I}^a)$, such that*

$$d\Omega^a = 0, \quad a = 1,\, 2,\, ...\, d, \tag{2.8}$$

$$\textstyle\sum_b \Omega^b = 0. \tag{2.9}$$

The vector space of Abelian relations of $\mathcal{W}$ is denoted by $\mathcal{A}(\mathcal{W})$. The Rank *of a Legendrian web $\mathcal{W}$ is the dimension of $\mathcal{A}(\mathcal{W})$.*

A Legendrian 1-web has rank zero by definition. Consider a Legendrian 2-web defined by a pair of sub-bundles $\mathcal{I}^a \subset T^*X$, $a = 1,\, 2$. Let $(\Omega^1,\, \Omega^2)$ be an Abelian relation. $\Omega^1 + \Omega^2 = 0$ and, since $\mathcal{I}^1 \cap \mathcal{I}^2 = \mathcal{C}$, Ω^1 must be a multiple of a contact 1-form. The defining equation $d\Omega^1 = 0$ then forces $\Omega^1 = \Omega^2 = 0$. Hence a Legendrian 2-web also has rank zero. Abelian relation has a meaning for Legendrian d-webs for $d \geq 3$.

The differential equation for Abelian relations (2.8), (2.9) is a linear differential system for the sections of the direct sum bundle $\oplus_{a=1}^d \mathcal{I}^a$. A computation for the case $d = 3,\, 4$ shows that it is over-determined, and one suspects that a generic Legendrian web has rank zero. The Legendrian webs we are interested in are those with many nontrivial Abelian relations. Moreover in this case, it turns out that the vector space $\mathcal{A}(\mathcal{W})$ inherits rich geometric structures from the layered feature of the integrability equations for (2.8), (2.9), Section 4.

In the opposite direction, it is not obvious if the rank of a Legendrian web is finite (for example, the characteristic variety argument in [3, p285] does not directly apply to Legendrian webs). One of the fundamental problems regarding the Abelian relation of a Legendrian web therefore would be to determine an effective bound on the

[5] In order the remove the ambiguity of adding by constants, we adopt here the differentiated version of first integrals, see the remark below Theorem 3.3.

rank, if such a bound exists. We shall present an argument in Section 4 that the expected maximum rank of a Legendrian d-web is $\rho_d = \frac{(d-1)(d-2)(2d+3)}{6}$.

The following example shows that the Legendrian webs with at least one Abelian relation exist in abundance.

Example 2.10. *Let X be a contact three manifold with the contact line bundle $\mathcal{C} \subset T^*X$. Let $\mathcal{W}_0$ be a Legendrian d-web ($d \geq 2$) defined by the d rank 2 sub-bundles $\mathcal{I}^a \subset T^*X$, $a = 1, 2, \dots d$. Let $\Omega^a \in C^\infty(\mathcal{I}^a)$ be a closed 1-form, $a = 1, 2, \dots d$. Assuming that Ω^a's are sufficiently generic, define another rank 2 sub-bundle $\mathcal{I}^{d+1} \subset T^*X$ by $\mathcal{I}^{d+1} = \mathcal{C} \oplus \langle \sum_b \Omega^b \rangle$. The $d+1$ rank 2 sub-bundles $\mathcal{I}^a \subset T^*X$, $a = 1, 2, \dots d+1$, define a Legendrian $(d+1)$-web which by construction has at least one nontrivial Abelian relation.*

3. Legendrian d-webs with $\rho_d = \frac{(d-1)(d-2)(2d+3)}{6}$ independent Abelian relations

In this section, we give an analysis of the Abelian relations for the Legendrian d-web $\mathcal{W}$ which is locally defined by the simple second order ODE's (3.1). An algebraic consideration leads to ρ_d linearly independent (un-differentiated) Abelian relations, each of whose components are polynomials in the natural adapted coordinate of $J^1(\mathbb{R}, \mathbb{R})$. The analysis in Section 4 will show that these account for all the Abelian relations of $\mathcal{W}$ and $\mathcal{W}$ has rank ρ_d.

Let $X = J^1(\mathbb{R}, \mathbb{R})$ equipped with the canonical contact structure, Example 1.4. Let (x, y, p) be the adapted coordinate so that the contact line bundle $\mathcal{C} = \langle dy - pdx \rangle$. Consider a Legendrian d-web $\mathcal{W}$ on X defined by the following set of second order ODE's;

$$y'' = q^a, \quad a = 1, 2, \dots d, \tag{3.1}$$
$$q^a\text{'s are distinct constants.}$$

The corresponding rank 2 sub-bundles of T^*X are defined by

$$\mathcal{I}^a = \langle\, dy - pdx,\ dp - q^a dx \,\rangle, \quad a = 1, 2, \dots d. \tag{3.2}$$

We claim that *$\mathcal{W}$ has at least ρ_d linearly independent Abelian relations.*[6]

Theorem 3.3. *Let $X = J^1(\mathbb{R}, \mathbb{R})$ be the 1-jet space of real valued functions on $\mathbb{R}$ equipped with the canonical contact structure. Let $\mathcal{W}$ be the Legendrian d-web on X defined by the second order ODE's* (3.1)*, $d \geq 3$. Then $\mathcal{W}$ has at least $\rho_d = \frac{(d-1)(d-2)(2d+3)}{6}$ linearly independent Abelian relations, and the rank of $\mathcal{W}$ is bounded below by ρ_d.*

The proof of theorem consists of the following four steps. In the first three steps, we construct ρ_d Abelian relations. In the last step, we show that they are linearly independent.

Let us indicate here a minor technical point for the analysis in this section. An Abelian relation for $\mathcal{W}$, (3.1), is by definition a d-tuple of closed 1-forms $(\Omega^1, \Omega^2, \dots \Omega^d)$, $\Omega^a \in C^\infty(\mathcal{I}^a)$. Fix a reference point $\mathrm{x}_0 = (0, 0, 0) \in X$, and let h^a be the unique local anti-derivative $dh^a = \Omega^a$ in a neighborhood of x_0 such that $h^a(\mathrm{x}_0) = 0$. It is clear that this defines an isomorphism from $\mathcal{A}(\mathcal{W})$ to the space of local un-differentiated Abelian relations, which consist of d-tuple of respective first integrals $(h^1, h^2, \dots h^d)$ vanishing at x_0 that satisfy the additive functional relation $\sum_b h^b = 0$. We shall freely use either of these equivalent forms as convenient.

Step 1. Universal first integrals

Consider the second order ODE

$$y'' = q, \tag{3.4}$$

where q is an indeterminate constant. We wish to construct the first integrals for (3.4) which are polynomials in q of the form

$$h(q) = h_0 + h_1\, q + h_2\, q^2 + \dots h_{m-1}\, q^{m-1}, \tag{3.5}$$

such that $h(q)|_{\mathrm{x}_0} = 0$.

[6] See Section 4 for the derivation of ρ_d.

Set the three first integrals for (3.4) that are linear in q;

$$u_0^3 = (y - p\,x) + q\,\frac{x^2}{2}, \tag{3.6}$$
$$u_1^3 = p - q\,x,$$
$$u_2^3 = \frac{p^2}{2} - q\,y.$$

Inductively define u_j^{m+1} for $m \geq 3,\ 0 \leq j \leq 2m-2$, by

$$u_j^{m+1} = u_0^3\,u_j^m, \qquad j = 0,\ 1,\ \dots 2m-4,$$
$$u_{2m-3}^{m+1} = u_2^3\,u_{2m-5}^m,$$
$$u_{2m-2}^{m+1} = u_2^3\,u_{2m-4}^m.$$

One has the formula

$$u_j^{m+1} = (u_0^3)^{j_0}\,(u_1^3)^{j_1}\,(u_2^3)^{j_2}, \quad 0 \leq j \leq 2m-2, \tag{3.7}$$

where the indices $(j_0,\ j_1,\ j_2)$ are uniquely determined by

$$j_0 + j_1 + j_2 = m-1, \tag{3.8}$$
$$j_0,\ j_2 \geq 0, \quad j_1 = 0, \text{ or } 1,$$
$$j = 0\cdot j_0 + 1\cdot j_1 + 2\cdot j_2.$$

Since these are polynomials in $\{\,u_0^3,\ u_1^3,\ u_2^3\,\}$, they are all first integrals for (3.4).

Assign the weights

$$\text{weight}(x,\ y,\ p,\ q) = (-1, 0, 1, 2)$$

respectively, and extend them to the polynomials in an obvious way. Let $\mathbb{R}[\,q\,]$ be the polynomial ring in the indeterminate q.

Lemma 3.9.

a) u_j^{m+1} *is homogenous of weight* j*, and the set of elements* $\{\,u_j^{m+1}\,\}_{j=0}^{2m-2}$ *are linearly independent over* $\mathbb{R}$ *as* $\mathbb{R}[\,q\,]$*-valued functions.*

b) u_j^{m+1} *has degree* $m-1$ *in* q*, and the two sets of elements* $\{\,u_j^{m+1}\,\}_{j=0}^{2m-2}$ *and* $\{\,u_j^{m'+1}\,\}_{j=0}^{2m'-2}$*,* $m \neq m'$*, are linearly independent over* $\mathbb{R}$ *as* $\mathbb{R}[\,q\,]$*-valued functions. Hence the set of elements* $\cup_{m=2}^{\infty}\{\,u_j^{m+1}\,\}_{j=0}^{2m-2}$ *are linearly independent over* $\mathbb{R}$ *as* $\mathbb{R}[\,q\,]$*-valued functions.*

Lemma follows from the construction of u_j^{m+1}. We shall omit the proof.

The following refined linear independence property will be used in **Step 4.**

Lemma 3.10. *Let* u *be a finite linear combination of the elements in* $\cup_{m=2}^{\infty}\{\,u_j^{m+1}\,\}_{j=0}^{2m-2}$*. If* $u(q_0) = 0$ *for a constant* q_0*, then* $u = 0$*.*

Proof. Note that $u_j^{m+1} \equiv y^{j_0}p^{j_1}(\frac{p^2}{2} - qy)^{j_2}, \mod x$, where $(j_0,\ j_1,\ j_2)$ is determined by (3.8). Evaluating at $q = q_0$, the highest weight term of $u_j^{m+1} \mod x$ (forgetting the weight of q) is $y^{j_0}p^{j_1+2j_2}$. From (3.8), since $j_1 = 0$, or 1, the associated index map $(m;\,j) \to (\,j_0,\ j = j_1 + 2j_2)$ is injective. Lemma follows from this by applying induction on decreasing weights. □

Step 2. Abelian relations

Let $\mathfrak{q}^l = ((q^1)^l,\ (q^2)^l,\ \dots (q^d)^l)$ denote a vector in $\mathbb{R}^d$ for $l = 0,\ 1,\ \dots d-2$. Vandermonde identity implies that there exists a linearly independent set of vectors $v^\mu = (v_1^\mu,\ v_2^\mu,\ \dots, v_d^\mu),\ \mu = 1,\ 2,\ \dots d-1$, such that

$$\langle\, v^\mu,\ \mathfrak{q}^l\,\rangle = \sum_a v_a^\mu\,(q^a)^l = 0, \quad \text{for } l = 0,\ 1,\ \dots d-\mu-1. \tag{3.11}$$

Take an element $h = u_j^{m+1},\ m \le d-1$, and consider the first integral $h(q^a)$ for $\mathcal{I}^a$, (3.2). By definition of u_j^{m+1}, it is a polynomial of the form $h(q^a) = h_0 + h_1\, q^a + h_2\, (q^a)^2 + \ldots, h_{m-1}\, (q^a)^{m-1}$, which is of degree $m-1$ in q. Then for each $v^\mu,\ 1 \le \mu \le d-m$, the d-tuple of first integrals

$$(v_1^\mu\, h(q^1),\ v_2^\mu\, h(q^2),\ \ldots,\ v_d^\mu\, h(q^d)) \tag{3.12}$$

gives an (un-differentiated) Abelian relation for $\mathcal{W}$ (the sum of the components vanish by (3.11)).

In summary, *each* u_j^{m+1}, *for* $2 \le m \le d-1$ *and* $0 \le j \le 2m-2$, *gives rise to* $d-m$ *Abelian relations for* $\mathcal{W}$.

Step 3. Decomposition of ρ_d

The preceding analysis suggests the following decomposition of ρ_d.

$$\begin{aligned}
\rho_3 :&\quad 1\cdot 3\\
\rho_4 :&\quad 2\cdot 3 + 1\cdot 5\\
\rho_5 :&\quad 3\cdot 3 + 2\cdot 5 + 1\cdot 7\\
\rho_6 :&\quad 4\cdot 3 + 3\cdot 5 + 2\cdot 7 + 1\cdot 9\\
&\quad \ldots\\
\rho_d :&\quad (d-2)\cdot 3 + (d-3)\cdot 5 + (d-4)\cdot 7 \quad \ldots \quad + 1\cdot(2d-3).
\end{aligned} \tag{3.13}$$

The set of universal first integrals $\{\, u_j^{m+1} \,\}_{j=0}^{2m-2}$ for $m = 2,\ 3,\ \ldots$, are identified with the diagonal entries $1\cdot 3,\ 1\cdot 5,\ 1\cdot 7,\ 1\cdot 9,\ \ldots$, in the above decomposition. The entries on the columns below the diagonal are identified with those generated by the appropriate vectors v^μ, (3.11), (3.12). One may check that this yields the correct formula for ρ_d.

Step 4. Linear independence

From **Step 2**, the given ρ_d Abelian relations are

$$\begin{aligned}
&\Gamma_j^{m;\,\mu} = (\, v_1^\mu\, u_j^{m+1}(q^1),\ v_2^\mu\, u_j^{m+1}(q^2),\ \ldots\ ,\ v_d^\mu\, u_j^{m+1}(q^d)\,),\\
&\text{for}\ \ 2 \le m \le d-1,\ \ 0 \le j \le 2m-2,\ \ 1 \le \mu \le d-m.
\end{aligned} \tag{3.14}$$

Suppose for a set of coefficients $c_{m;\,\mu}^j$ a linear combination $\sum_{m;\,\mu,\,j} c_{m;\,\mu}^j\, \Gamma_j^{m;\,\mu} = 0$. For each $a = 1,\ 2,\ \ldots d$,

$$\sum_{m;\,j} \Big(\, \sum_\mu c_{m;\,\mu}^j v_a^\mu \Big)\, u_j^{m+1}(q^a) = 0.$$

By Lemma 3.10, this implies $\sum_{m;\,j} \big(\, \sum_\mu c_{m;\,\mu}^j v_a^\mu \big)\, u_j^{m+1} = 0,\ a = 1,\ 2,\ \ldots d$. By Lemma 3.9, this implies for each $m;\ j$ that $\sum_\mu c_{m;\,\mu}^j v_a^\mu = 0,\ a = 1,\ 2,\ \ldots d$. The linear independence of the vectors $v^\mu = (v_1^\mu,\ v_2^\mu,\ \ldots v_d^\mu)$, (3.11), then forces $c_{m;\,\mu}^j = 0$. $\square$

4. Maximum rank of a Legendrian web

In the last section, we gave an algebraic construction of the Abelian relations for a particular class of Legendrian webs. In this section, we give an analysis of the linear differential system for Abelian relations of the general Legendrian webs. As a corollary, we show that the rank of a Legendrian d-web is bounded by ρ_d.

The idea is to successively differentiate the defining equations (2.8), (2.9) for Abelian relations, and to analyze the higher order compatibility equations. We present two different analysis of these compatibility equations. The first analysis shows that the linear differential system for Abelian relations of a Legendrian d-web closes up at weight $d-2$, provided that an associated sequence of universal integer coefficient matrices are non-degenerate in a strict sense, Theorem 4.24. The second analysis shows that this linear differential system closes up at depth $2d-4$, and the rank is bounded by ρ_d, Theorem 4.37. See Definition 4.8 for the definition of weight, and depth.

Let us give an outline of the analysis. Given a Legendrian d-web, we first determine the infinitely prolonged structure equation for the first integrals of each Legendrian foliation, Lemma 4.12. The structure equation is applied to the defining additive equation of Abelian relation, and one obtains a sequence of higher order compatibility equations, Proposition 4.17. The prolonged structure equation, and the sequence of compatibility equations lead to the three filtrations of the infinite jets of an Abelian relation by *weight*,[7] *height*, and *depth*. An examination of the compatibility equations according to this triple filtrations shows that the aforementioned sequence of integer coefficient matrices represents in a way a part of the symbol of the prolonged linear differential system for Abelian relations, and Theorem 4.24 follows from the general theory of over-determined PDE. The rank bound in Theorem 4.37 is obtained by examining the symbol as a whole with respect to the known Abelian relations from Section 3.

Throughout this section, we shall not use the summation convention on repeated indices, and instead use the summation sign.

4.1. **Structure equation.**

4.1.1. *Prolongation.* We continue the analysis from Section 2.1.

Let $\mathcal{W}$ be a Legendrian d-web on a contact three-manifold X. Let (x, y, p) be an adapted local coordinate of X so that $\mathcal{W}$ is defined by the d second order ODE's (2.2). The corresponding rank 2 sub-bundles of T^*X are

$$\mathcal{I}^a = \langle\, dy - pdz,\ dp - q^a(x, y, p)dx \,\rangle, \quad a = 1, 2, \dots d, \tag{4.1}$$

where $q^a \neq q^b$ for $a \neq b$.

Set

$$\begin{aligned} \theta &= dy - p\, dx, \\ \theta^a &= dp - q^a\, dx, \quad a = 1,\ 2,\ \dots d, \end{aligned} \tag{4.2}$$

and let

$$\Omega^a = f^a_{10}\theta^a + f^a_{01}\theta, \quad a = 1,\ 2,\ \dots d, \tag{4.3}$$

denote a section of $\mathcal{I}^a$ for coefficients $\{\, f^a_{10},\ f^a_{01} \,\}_{a=1}^d$.

The vector space of Abelian relations $\mathcal{A}(\mathcal{W})$ is by definition the space of solutions to the linear differential system (2.8), (2.9). It is written as a system of first order linear differential equations for the coefficients $\{\, f^a_{10},\ f^a_{01} \,\}_{a=1}^d$ as follows.

$$(df^a_{10} + (f^a_{01} + q^a_p f^a_{10})dx) \wedge \theta^a + (df^a_{01} + q^a_y f^a_{10} dx) \wedge \theta = 0, \quad a = 1,\ 2,\ \dots d, \tag{4.4}$$

$$\begin{aligned} &\textstyle\sum f^b_{10} = 0,\ \ \sum q^b f^b_{10} = 0, \\ &\textstyle\sum f^b_{01} = 0. \end{aligned} \tag{4.5}$$

Here $q^a_p,\ q^a_y$ denote the partial derivatives.

We shall determine the infinite prolongation of (4.4). By Cartan's lemma, there exist the derivative coefficients $\{\, f^a_{20},\ f^a_{11},\ f^a_{02} \,\}_{a=1}^d$ such that

$$\begin{aligned} df^a_{10} &= f^a_{20}\theta^a + f^a_{11}\,\theta - (\, f^a_{01} + \tau^a_{10})\, dx, \\ df^a_{01} &= f^a_{11}\theta^a + f^a_{02}\,\theta - (\qquad\ \ \tau^a_{01})\, dx, \quad a = 1,\ 2,\ \dots d, \end{aligned} \tag{4.6}$$

where $\tau^a_{10} = q^a_p f^a_{10},\ \tau^a_{01} = q^a_y f^a_{10}$. Note the identity $\tau^a_{10},\ \tau^a_{01} \equiv 0 \mod f^a_{10}$.

By Cartan's lemma again (or equivalently since mixed partials commute), it is clear that one may inductively define $\{\, f^a_{ij} \,\}_{a=1}^d$ for $i,\ j \geq 0,\ (i\, j) \neq (0\, 0)$, by

$$df^a_{ij} \equiv f^a_{(i+1)j}\theta^a + f^a_{i(j+1)}\,\theta, \mod dx. \tag{4.7}$$

Note that

$$f^a_{(i+1)j} = \frac{\partial}{\partial p} f^a_{ij}, \quad f^a_{i(j+1)} = \frac{\partial}{\partial y} f^a_{ij}.$$

[7] The 'weight' in this section is different from the weight used in Section 3.




Figure 4.1. Υ_{ij} for the case $(i\,j) = (3\,1)$: upper-left chamber.
A node with (ij) on top represents the set $\{ f^a_{ij} \}_{a=1}^d$. The weight increases as one moves to the right, the height increases as one moves upward, and the depth increases as one moves to the lower-right. The chamber Υ_{ij} consists of the $f^a_{i'j'}$'s represented by the node $(i'j')$ which lies in the upper-left chamber bounded by the vertical weight line, and the slant depth line.

Set $f^a_{ij} = 0$ for $i < 0,\ j < 0$, or $(i\,j) = (0\,0)$.

In order to facilitate the computation, let us introduce the following three filtrations for the coefficients f^a_{ij}'s.

Definition 4.8. *Let* $\{ f^a_{ij} \}_{a=1}^d;\ \ i,\, j \ \geq\ 0, (i\,j) \ \neq\ (0\,0)$, *be the successive derivatives of the coefficients* $\{ f^a_{10},\, f^a_{01} \}_{a=1}^d$ *for an Abelian relation* (4.3), (4.4), (4.5). *Assign*

$$\begin{aligned} \text{height}(f^a_{ij}) &= i, \\ \text{weight}(f^a_{ij}) &= i + j, \\ \text{depth}(f^a_{ij}) &= i + 2j. \end{aligned} \tag{4.9}$$

For a finite linear combination of f^a_{ij}*'s, weight(depth) is defined as the maximum weight(depth) of nonzero terms.*

The weight(depth) of an Abelian relation (4.3) *(or more generally a first integral) is the maximum weight(depth) of the associated nonzero derivatives* f^a_{ij}.

The fundamental chamber Υ_{ij} at $(i\,j)$ *is the subset*

$$\Upsilon_{ij} = \{\, f^a_{i'j'} \mid\ \text{weight}(f^a_{i'j'}) \leq i + j,\ \ \text{depth}(f^a_{i'j'}) \leq i + 2j \ \}. \tag{4.10}$$

The subset of elements $\{ f^a_{ij} \}_{a=1}^d \subset \Upsilon_{ij}$ are characterized as the *dominant* coefficients in that they are exactly the elements for which the equalities hold in the definition of Υ_{ij}. Note by definition the relation (with an abuse of notation)

$$\begin{aligned} \frac{\partial}{\partial p}\Upsilon_{ij} &\subset \Upsilon_{(i+1)j}, \\ \frac{\partial}{\partial y}\Upsilon_{ij} &\subset \Upsilon_{i(j+1)}, \\ (\frac{\partial}{\partial x} + p\frac{\partial}{\partial y})\Upsilon_{ij} &\subset \Upsilon_{(i+1)j}. \end{aligned} \tag{4.11}$$

(this is obvious except the third one, (4.22)). This inclusion relation will be used implicitly for the rest of this section.

The following refinement of (4.7) suffices for our purpose.

Lemma 4.12. *The rough structure equation* (4.7) *can be refined to*

$$df^a_{ij} = f^a_{(i+1)j}\theta^a + f^a_{i(j+1)}\,\theta - (\,i f^a_{(i-1)(j+1)} + \tau^a_{ij}\,)\,dx, \tag{4.13}$$

where

$$\tau^a_{ij} \equiv 0, \quad \mod\ \Upsilon_{ij}. \tag{4.14}$$

Proof. We apply induction argument on increasing depth, and weight.

Differentiating (4.13) by the Lie derivative $L_{\frac{\partial}{\partial p}}$, one gets

$$\begin{aligned} df^a_{(i+1)j} = {} & f^a_{(i+2)j}\theta^a + f^a_{(i+1)(j+1)}\,\theta - (i\, f^a_{i(j+1)} + L_{\frac{\partial}{\partial p}}\tau^a_{ij})\,dx \\ & - (f^a_{(i+1)j} q^a_p + f^a_{i(j+1)})\,dx. \end{aligned}$$

By the induction hypothesis and (4.11), $\{\, f^a_{(i+1)j},\, L_{\frac{\partial}{\partial p}}\tau^a_{ij}\,\} \subset \Upsilon_{(i+1)j}$. Hence $\tau^a_{(i+1)j} \equiv 0, \mod\ \Upsilon_{(i+1)j}$.

Differentiating (4.13) by the Lie derivative $L_{\frac{\partial}{\partial y}}$, one gets

$$\begin{aligned} df^a_{i(j+1)} = {} & f^a_{(i+1)(j+1)}\theta^a + f^a_{i(j+2)}\,\theta - (i\, f^a_{(i-1)(j+2)} + L_{\frac{\partial}{\partial y}}\tau^a_{ij})\,dx \\ & - f^a_{(i+1)j} q^a_y\,dx. \end{aligned}$$

By the induction hypothesis, one gets $\tau^a_{i(j+1)} \equiv 0, \mod\ \Upsilon_{i(j+1)}$.

(4.14) now follows from the initial condition (4.6) that $\tau^a_{10}, \tau^a_{01} \equiv 0, \mod\ \Upsilon_{10}$. □

4.1.2. *Compatibility equations.* In this sub-section, we apply Lemma 4.12 repeatedly to compute the successive derivatives of the additive equations (4.5).

There are two preliminaries. Firstly, let us introduce a simplified notation

$$\langle \mathfrak{q}^i,\, f_{jk}\rangle = \sum_b (q^b)^i f^b_{jk}.$$

The defining equations (4.5) are written as

$$\begin{aligned} &\langle \mathfrak{q}^0,\, f_{10}\rangle = 0,\ \ \langle \mathfrak{q}^1,\, f_{10}\rangle = 0, \\ &\langle \mathfrak{q}^0,\, f_{01}\rangle = 0. \end{aligned} \tag{4.15}$$

Secondly, set an ∞-by-∞ integer matrix (c^I_J) for $I \geq 0$ as follows.

$$\begin{aligned} & c^I_0 = 1, \\ & c^I_J = 0, \qquad \text{for } J < 0, \text{ or } J > \frac{I}{2}, \\ & c^I_J = (I - 2J + 1)\, c^{I-1}_{J-1} + c^{I-1}_J, \\ & \quad = \frac{I!}{2^J\,(I-2\,J)!\,J!}. \end{aligned} \tag{4.16}$$

The first few terms are (for $J = 0,\, 1,\, 2,\, 3$),

c^0_J	1	0	0	0
c^1_J	1	0	0	0
c^2_J	1	1	0	0
c^3_J	1	3	0	0
c^4_J	1	6	3	0

Lemma 4.17. *The successive derivatives of the defining equations* (4.5) *for the Abelian relations via Lemma 4.12 yield the following sequence of higher order compatibility equations.*

$$Eq^I_{ij}: \quad c^I_0 \langle \mathfrak{q}^I, f_{ij}\rangle + c^I_1 \langle \mathfrak{q}^{I-1}, f_{(i-2)(j+1)}\rangle \dots + c^I_k \langle \mathfrak{q}^{I-k}, f_{(i-2k)(j+k)}\rangle \dots \equiv 0, \quad \mod \Upsilon_{(i+1)(j-1)}, \qquad \text{for } 0 \le I \le i. \tag{4.18}$$

Proof. We apply induction argument on increasing depth, and weight.

Step 1. Differentiating (4.18) by the Lie derivative $L_{\frac{\partial}{\partial y}}$, and noting that $L_{\frac{\partial}{\partial y}} \Upsilon_{(i+1)(j-1)} \subset \Upsilon_{(i+1)j}$, one gets

$$Eq^I_{i(j+1)}: \quad c^I_0 \langle \mathfrak{q}^I, f_{i(j+1)}\rangle + c^I_1 \langle \mathfrak{q}^{I-1}, f_{(i-2)(j+2)}\rangle \dots + c^I_k \langle \mathfrak{q}^{I-k}, f_{(i-2k)(j+k+1)}\rangle \dots \equiv 0, \quad \mod \Upsilon_{(i+1)j}, \qquad \text{for } 0 \le I \le i.$$

It thus suffices to verify the case $j = 0$.

Step 2. Differentiating (4.18) by the Lie derivative $L_{\frac{\partial}{\partial p}}$, and noting that $L_{\frac{\partial}{\partial p}} \Upsilon_{(i+1)(j-1)} \subset \Upsilon_{(i+2)(j-1)}$, one gets

$$Eq^I_{(i+1)j}: \quad c^I_0 \langle \mathfrak{q}^I, f_{(i+1)j}\rangle + c^I_1 \langle \mathfrak{q}^{I-1}, f_{(i-1)(j+1)}\rangle \dots + c^I_k \langle \mathfrak{q}^{I-k}, f_{(i-2k+1)(j+k)}\rangle \dots \equiv 0, \quad \mod \Upsilon_{(i+2)(j-1)}, \qquad \text{for } 0 \le I \le i.$$

It thus suffices to verify the case $I = i$.

Step 3. Differentiating (4.18) for the case $I = i$ by the Lie derivative $L_{\frac{\partial}{\partial x}+p\frac{\partial}{\partial y}}$, and noting that $L_{\frac{\partial}{\partial x}+p\frac{\partial}{\partial y}} \Upsilon_{(i+1)(j-1)} \subset \Upsilon_{(i+2)(j-1)}$, one gets from (4.22)

$$\begin{aligned} Eq^{i+1}_{(i+1)j}: \quad & c^i_0 \left(\langle \mathfrak{q}^{i+1}, f_{(i+1)j}\rangle + i \langle \mathfrak{q}^i, f_{(i-1)(j+1)}\rangle\right) + c^i_1 \left(\langle \mathfrak{q}^i, f_{(i-1)(j+1)}\rangle + (i-2)\langle \mathfrak{q}^{i-1}, f_{(i-3)(j+2)}\rangle\right) \\ & \dots c^i_k \left(\langle \mathfrak{q}^{i+1-k}, f_{(i+1-2k)(j+k)}\rangle + (i-2k)\langle \mathfrak{q}^{i-k}, f_{(i-2k-1)(j+k+1)}\rangle\right) \dots \equiv 0, \\ & \mod \Upsilon_{(i+2)(j-1)}. \end{aligned}$$

This implies the recursive relation

$$\begin{aligned} c^{i+1}_0 &= c^i_0, \\ c^{i+1}_1 &= i\, c^i_0 + c^i_1, \\ &\dots \\ c^{i+1}_k &= (i-2k+2)\, c^i_{k-1} + c^i_k, \\ &\dots . \end{aligned}$$

(4.18) follows from the initial condition (4.15). □

In summary, Lemma 4.17 shows that;

• as one differentiates, the compatibility equations for Abelian relations propagate in the direction of increasing depths,

• for each fundamental chamber Υ_{ij}, there are $(i+1)$ linear compatibility equations for the dominant coefficients $\{ f^a_{ij} \}^d_{a=1}$.

In the next section, we shall give a geometric interpretation of these results.

4.2. **Rank bound - an analysis.** In this section, the compatibility equations (4.18) are combined in a sequence of appropriately layered manner to show that the proof of rank bound can be reduced to the non-degeneracy of a sequence of integer coefficient matrices.

4.2.1. *Overview.* Let us first explain our claim. Recall the definition of weight$(f^a_{ij}) = i + j$, Definition 4.8. Set

$$F_w = \{ f^a_{ij} \mid \text{weight}(f^a_{ij}) = w \}, \tag{4.19}$$
$$\mathcal{F}_w = \{ f^a_{ij} \mid \text{weight}(f^a_{ij}) \leq w \}.$$

Our claim in this section is that (symbolically)

$$d\mathcal{F}_{d-2} \equiv 0, \mod \mathcal{F}_{d-2}, \tag{4.20}$$

and *the linear differential system for the Abelian relations closes up at weight* $d-2$.

Assuming this claim is true, the rank bound is obtained by counting the number of linearly independent elements in $\mathcal{F}_{d-2}$. Since $F_w = \{ f^a_{w0}, f^a_{(w-1)1}, \dots f^a_{0w} \}^d_{a=1}$, (4.18) implies that, modulo lower weight terms, there are $(d-(w+1)) + (d-w) + \dots (d-1)$ linearly independent elements. The rank bound is given by

$$\sum_{w=1}^{d-2} (d-(w+1)) + (d-w) + \dots (d-1) = \rho_d.$$

Let us point out a possibly unusual phenomenon that this claim entails. In the theory of closed, over-determined differential systems, the number of admissible derivatives generally decreases as one differentiates, until it closes up at certain order and all the higher order derivatives are expressed as functions of the derivatives up to that order. On the contrary in our case the number of linearly independent elements in F_w increases until the differential system suddenly closes up at weight $d-2$.

One may explain this by using the filtration by depth rather than by weight. The vector space of Abelian relations $\mathcal{A}(\mathcal{W})$ is by definition a section of the direct sum bundle $\oplus^d_{a=1}\mathcal{I}^a$. Consider the projection

$$\mathcal{A}(\mathcal{W}) \hookrightarrow C^\infty(\oplus^d_{a=1}\mathcal{I}^a) \to C^\infty(\oplus^d_{a=1}\mathcal{I}^a/\mathcal{C}),$$

where $\mathcal{C}$ is the contact line bundle. Since the components of each elements of $\mathcal{A}(\mathcal{W})$ is a closed 1-form, the projection map is injective. This implies that, in order to determine a rank bound, it suffices to analyze the higher order derivatives modulo $\mathcal{C}$. In hindsight, this is equivalent to the following elementary fact.[8]

Let X be a contact three manifold. Let $\mathcal{C} \subset T^*X$ be the contact line bundle. Let $Z = (Z_1, Z_2, \dots Z_r)$ be an $\mathbb{R}^r$-valued function on X (here r stands for the rank of $\mathcal{W}$). Suppose it satisfies the equation

$$dZ \equiv 0, \mod Z; \mathcal{C}.$$

Then in fact

$$dZ \equiv 0, \mod Z. \tag{4.21}$$

The structure equation in Lemma 4.12 gives a more concrete description. Since differentiating by $\frac{\partial}{\partial p}$, and $\frac{\partial}{\partial x} + p\frac{\partial}{\partial y}$ is equivalent to differentiating modulo $\mathcal{C}$, one notes the identity

$$\frac{\partial}{\partial p} f^a_{ij} \equiv f^a_{(i+1)j}, \tag{4.22}$$
$$(\frac{\partial}{\partial x} + p\frac{\partial}{\partial y}) f^a_{ij} \equiv -q^a f^a_{(i+1)j} - i f^a_{(i-1)(j+1)}, \qquad \mod \Upsilon_{ij}.$$

The differential operators $\frac{\partial}{\partial p}$, $\frac{\partial}{\partial x} + p\frac{\partial}{\partial y}$ increase the depth by 1 at most.

Returning to counting the number of linearly independent elements in $\mathcal{F}_{d-2}$, suppose we filter $\mathcal{F}_{d-2}$ by depths. Then, as one may visualize in the diagram Fig. 4.1, the number of linearly independent elements of depth δ generally increases up to $\delta = d-2$. At this depth, according to our claim, the number of linearly independent elements starts to decrease, until $\delta = 2(d-2)+2$ and the linear differential system closes up.

[8]This observation is the basis of the proof of rank bound in Section 4.3.

4.2.2. *Sub-symbol.* In order to state the main result of this sub-section, there are two preliminaries.

Firstly, the analysis of the compatibility equations (4.18) requires manipulating the matrices that consist of the blocks of Vandermonde-like matrices. Let us introduce a simplified notation for these. Recall the definition of c^I_J, (4.16). Let $\mathfrak{q}^I = ((q^1)^I, (q^2)^I, \dots (q^d)^I)$ denote the row vector. Let $V_J^{I_1, I_2}$ be the $(I_2 - I_1 + 1)$-by-d matrix

$$V_J^{I_1, I_2} = \begin{pmatrix} c_J^{I_1}\, \mathfrak{q}^{I_1 - J} \\ c_J^{I_1+1}\, \mathfrak{q}^{I_1+1-J} \\ \dots \\ c_J^{I_2}\, \mathfrak{q}^{I_2 - J} \end{pmatrix}.$$

For example, $V_0^{0,\, d-1}$ is a standard Vandermonde matrix of size d.

Secondly, the analysis will show that the condition for such a generalized Vandermonde matrix to be non-degenerate is equivalent to that a set of matrices with c^I_J's as components are non-degenerate in a strict sense.

Definition 4.23. *A square matrix is* strongly non-degenerate *when each of its upper-left corner square sub-matrices, including the matrix itself, is non-degenerate.*

The typical example of a strongly non-degenerate matrix is a Vandermonde matrix.

Let us now state the main result.

Theorem 4.24. *Let c^I_J be the integer sequence defined by* (4.16)*. The linear differential system* (4.4), (4.5) *for Abelian relations of a Legendrian d-web on a contact three manifold closes up at weight $d-2$, and the rank is bounded above by $\rho_d = \frac{(d-1)(d-2)(2\,d+3)}{6}$, provided that the following set of matrices are strongly non-degenerate.*

$$\begin{pmatrix} c_1^d & c_2^d & \dots & c_{d-2}^d & c_{d-1}^d \\ c_0^{d-1} & c_1^{d-1} & \dots & c_{d-3}^{d-1} & c_{d-2}^{d-1} \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & c_0^3 & c_1^3 & c_2^3 \\ \dots & \dots & \dots & c_0^2 & c_1^2 \end{pmatrix}, \tag{4.25}$$

$$\begin{pmatrix} c_2^{d+1} & c_3^{d+1} & \dots & c_{d-1}^{d+1} \\ c_1^d & c_2^d & \dots & c_{d-2}^d \\ \dots & \dots & \dots & \dots \\ \dots & \dots & c_1^4 & c_2^4 \end{pmatrix},$$

$$\dots$$

$$\begin{pmatrix} c_{d-2}^{2d-3} & c_{d-1}^{2d-3} \\ c_{d-3}^{2d-4} & c_{d-2}^{2d-4} \end{pmatrix},$$

$$c_{d-1}^{2d-2}.$$

Proof. Throughout the proof, it is helpful to consult the diagram Fig. 4.1.

Recall the notations in Section 4.2.1. The claim is that $F_{d-1} = \{\, f^a_{(d-1)0},\, f^a_{(d-2)1},\, \dots f^a_{0(d-1)} \,\}_{a=1}^d$ is determined by $\mathcal{F}_{d-2}$ via the compatibility equations (4.18). We shall apply an induction on increasing depth.

Depth d-1. Consider $\{\, f^a_{(d-1)0} \,\}_{a=1}^d$. (4.18) implies that they satisfy d equations

$$\langle\, c_0^I\, \mathfrak{q}^I,\, f_{(d-1)0} \,\rangle \equiv 0, \quad \mod \mathcal{F}_{d-2}\, (\text{since } \Upsilon_{d(-1)} \subset \mathcal{F}_{d-2}), \quad 0 \le I \le d-1. \tag{4.26}$$

Taking $f_{(d-1)0} = (\, f^1_{(d-1)0},\, f^2_{(d-1)0},\, \dots f^d_{(d-1)0}\,)^t$ as a column vector, the claim is that the d-by-d matrix $V_0^{0,d-1}$ is non-degenerate. $c_0^I = 1$, and $V_0^{0,d-1}$ is a Vandermonde matrix, and hence non-degenerate.

Depth d. Consider $\{ f^a_{d0},\, f^a_{(d-2)1} \}_{a=1}^d$. (4.18) implies that they satisfy $2d$ equations

$$\begin{aligned} \langle\, c_0^I \mathfrak{q}^I,\, f_{d0} \,\rangle + \langle\, c_1^I \mathfrak{q}^{I-1},\, f_{(d-2)1} \,\rangle &\equiv 0, \quad \mod\ \mathcal{F}_{d-2}, \quad 0 \le I \le d, \\ &\quad (\text{since } \Upsilon_{(d+1)(-1)} \subset \mathcal{F}_{d-2} \text{ by } \textbf{Depth d-1}), \\ \langle\, c_0^I \mathfrak{q}^I,\, f_{(d-2)1} \,\rangle &\equiv 0, \quad \mod\ \mathcal{F}_{d-2}, \quad 0 \le I \le d-2, \\ &\quad (\text{since } \Upsilon_{(d-1)0} \subset \mathcal{F}_{d-2} \text{ by } \textbf{Depth d-1}). \end{aligned}$$

After some row changes, the claim is that the following $2d$-by-$2d$ matrix is non-degenerate.

$$\begin{pmatrix} V_0^{0,d-1} & V_1^{0,d-1} \\ . & V_0^{0,d-2} \\ V_0^{d,d} & V_1^{d,d} \end{pmatrix},$$

where $'.'$ denotes 0 matrix.

An elementary row operation shows that this matrix is non-degenerate when

$$c_1^d \neq 0. \tag{4.27}$$

We assume this from now on.

Depth d+1. Arguing similarly as above, one may verify that, after some row changes, the claim is equivalent to that the following $3d$-by-$3d$ matrix is non-degenerate.

$$\begin{pmatrix} V_0^{0,d-1} & V_1^{0,d-1} & V_2^{0,d-1} \\ . & V_0^{0,d-1} & V_1^{0,d-1} \\ . & . & V_0^{0,d-3} \\ V_0^{d,d+1} & V_1^{d,d+1} & V_2^{d,d+1} \end{pmatrix}.$$

The d-by-d sub-matrix $V_0^{0,d-1}$ in the center is non-degenerate by **Depth d-1**. Elementary row operations show that this matrix is non-degenerate when

$$\begin{aligned} \det \begin{pmatrix} c_1^d & c_2^d \\ c_0^{d-1} & c_1^{d-1} \end{pmatrix} &\neq 0, \\ c_2^{d+1} &\neq 0. \end{aligned} \tag{4.28}$$

Note that the previous non-degeneracy condition (4.27) can be combined with (4.28) to imply that the two matrices in (4.28) are strongly non-degenerate. We assume this from now on.

Depth d+2. Arguing similarly as above, one may verify that, after some row changes, the claim is equivalent to that the following $4d$-by-$4d$ matrix is non-degenerate.

$$\begin{pmatrix} V_0^{0,d-1} & V_1^{0,d-1} & V_2^{0,d-1} & V_3^{0,d-1} \\ . & V_0^{0,d-1} & V_1^{0,d-1} & V_2^{0,d-1} \\ . & . & V_0^{0,d-2} & V_1^{0,d-2} \\ . & V_0^{d,d} & V_1^{d,d} & V_2^{d,d} \\ . & . & . & V_0^{0,d-4} \\ V_0^{d,d+2} & V_1^{d,d+2} & V_2^{d,d+2} & V_3^{d,d+2} \end{pmatrix}.$$




Figure 4.2. Node diagram for the case $d = 4$.

The $2d$-by-$2d$ sub-matrix in the center is non-degenerate by **Depth d**. Elementary row operations show that this matrix is non-degenerate when

$$\det \begin{pmatrix} c_1^d & c_2^d & c_3^d \\ c_0^{d-1} & c_1^{d-1} & c_2^{d-1} \\ c_{-1}^{d-2} & c_0^{d-2} & c_1^{d-2} \end{pmatrix} \neq 0, \tag{4.29}$$

$$\det \begin{pmatrix} c_2^{d+1} & c_3^{d+1} \\ c_1^d & c_2^d \end{pmatrix} \neq 0,$$

$$c_3^{d+2} \neq 0.$$

Note that the previous non-degeneracy conditions (4.27), (4.28) can be combined with (4.29) to imply that the three matrices in (4.29) are strongly non-degenerate. We assume this from now on.

Induction argument. Proceeding in this way using induction on increasing odd or even depths up to **Depth 2d-2**, the strong non-degeneracy of the $d-1$ matrices (4.25) implies that

$$F_{d-1} \equiv 0, \mod \mathcal{F}_{d-2}. \tag{4.30}$$

The linear differential system for Abelian relations is then closed at weight $d-2$. $\square$

We are currently unable to prove that the matrices (4.25) are strongly non-degenerate for all $d \geq 3$.

Remark 4.31. *The strong non-degeneracy of the set of matrices* (4.25) *may be checked for small values of* d *on the computer. It is true for* d *from 3 up to 100. The* `Maple` *code for this computation is included in Appendix.*

Example 4.32. *Consider the case* $d = 4$*, Fig. 4.2. Each '*$\bullet$*' represents the elements of* $\mathcal{F}_{d-2} = \mathcal{F}_2$*. Each '*$\otimes$*' represents the elements of* $F_{d-1} = F_3$*, which is to be determined by* $\mathcal{F}_2$*. Each '*$\circ$*' on a slant line represents the coefficients that give complementary linear equations for the lowest end* $\otimes$*.*

The thick horizontal line of height $d-1=3$ *bisects the slant lines, which implies that the number of linear compatibility equations match the number of unknowns represented by the nodes* $\otimes$ *and* $\circ$ *connected by a slant line. The analysis in this section shows that this system of linear equations has full rank for each depth from depth* $d-1=3$ *to depth* $2d-2=6$*, provided that the integer matrices* (4.25) *for* $d=4$ *are strongly non-degenerate.*

4.3. **Proof of rank bound.** Theorem 4.24 makes use of a part of the symbol of the linear differential system for Abelian relations. It specifies the subset of variables f^a_{ij} which span the remaining variables, and attempts to show that the differential system closes up at weight $d-2$.

In this section, we utilize all of the symbol. We show that the linear differential system for Abelian relations closes up at depth $2d-4$, and the rank of a Legendrian d-web is bounded by the expected value ρ_d.

Recall the definition of depth$(f^a_{ij}) = i+2j$, Definition 4.8. Set

$$\mathcal{F}^\delta = \{\, f^a_{ij} \mid \text{depth}(f^a_{ij}) \le \delta \,\}.$$

Our claim in this section is that

$$d\mathcal{F}^{2d-4} \equiv 0, \quad \mod \ \mathcal{F}^{2d-4}, \tag{4.33}$$

and *the linear differential system for Abelian relations closes up at depth* $2d-4$.

Consider the linear differential system for Abelian relations of depth $\le (2d-4)+1$ of the Legendrian webs considered in Section 3.

$$\begin{aligned} df^a_{ij} &= f^a_{(i+1)j}\theta^a + f^a_{i(j+1)}\,\theta - \, i f^a_{(i-1)(j+1)}\,dx, \\ f^a_{ij} &= 0 \ \text{ for } \ (i\,j) = (0\,0), \ i<0, \ j<0, \ i+2j > (2d-4)+1, \end{aligned} \tag{4.34}$$

with the linear constraints

$$\begin{aligned} Eq^I_{ij}: \quad & c^I_0 \langle \mathfrak{q}^I, f_{ij}\rangle + c^I_1 \langle \mathfrak{q}^{I-1}, f_{(i-2)(j+1)}\rangle \,... \, + c^I_k \langle \mathfrak{q}^{I-k}, f_{(i-2k)(j+k)}\rangle \,... \, = 0, \\ & \text{for } \ 0 \le I \le i. \end{aligned} \tag{4.35}$$

Here $\{\, \theta^a = dp - q^a dx,\, \theta = dy - p dx \,\}$; $\mathfrak{q}^I = ((q^1)^I,\, (q^2)^I,\, ...\, (q^d)^I)^t$, $f_{ij} = (f^1_{ij},\, f^2_{ij},\, ...\, f^d_{ij})^t$, and q^a's are distinct constants. This system of equations is compatible by the arguments in the proof of Lemma 4.12, Lemma 4.17.

For each depth $1,\, 2,\, 3,\, ...$, one has

depth	number of variables f^a_{ij}	number of compatibility equations E^I_{ij}
1	d	2
2	$2d$	$1+3$
3	$2d$	$2+4$
4	$3d$	$1+3+5$
5	$3d$	$2+4+6$
...	...	...
	kd	k^2
	kd	$k(k+1)$
...	...	...
$2d-4$	$(d-1)d$	$(d-1)^2$
$2d-3$	$(d-1)d$	$(d-1)d$

The total sum is

$$\begin{aligned} &(\text{ number of variables }) \ - \ (\text{ number of compatibility equations }) \\ &= \left(d + \sum_{k=2}^{d-1} 2kd \right) - \left(2 + \sum_{k=2}^{d-1} k^2 + k(k+1) \right) = \rho_d. \end{aligned} \tag{4.36}$$

The analysis of Section 4.2 implies that our claim (4.33) is equivalent to that the $(d-1)d$ constraint equations of depth $2d-3$ are linearly independent. We shall give a proof by the following argument. Let $\mathcal{S}_d$ denote the vector space of solutions to (4.34) with (4.35). Since this linear differential system is compatible, (4.36) implies

that $\dim \mathcal{S}_d$ is at least ρ_d, and is strictly larger than ρ_d whenever the given set of constraint equations are linearly dependent. It thus suffices to show that; a) $\dim \mathcal{S}_d \leq \rho_d$, b) every solution in $\mathcal{S}_d$ has depth $\leq 2d-4$.

Theorem 4.37. *For $d \geq 3$, the rank of a Legendrian d-web is bounded above by $\rho_d = \frac{(d-1)(d-2)(2d+3)}{6}$.*

Theorem 4.37 now follows from the following refinement of Theorem 3.3.

Lemma 4.38. *Let $\mathcal{W}$ be the Legendrian d-web defined by the second order ODE's* (3.1)*, $d \geq 3$. The set of ρ_d Abelian relations* (3.14) *are exactly the set of Abelian relations of depth $\leq (2d-4)+1$, which are necessarily of depth $\leq 2d-4$.*

Proof. **Step 1**. An Abelian relation of depth $\leq (2d-4)+1$ has weight $\leq (2d-4)+1$ by definition of depth and weight. Since the weight equals the degree as a polynomial in the variables $\{p, y\}$, an Abelian relation of depth $\leq (2d-4)+1$ is a polynomial in the variables $\{p, y\}$.

The first integrals $\{ u_0^3(q^a), u_1^3(q^a) \}$ for the foliation defined by the second order ODE $y'' = q^a$ are functionally independent (note $u_0^3(q^a) \equiv y,\ u_1^3(q^a) \equiv p, \mod x$). One may in fact linearly solve for the variables $\{ p, y \}$ in terms of $\{ u_0^3(q^a), u_1^3(q^a) \}$. A first integral for the given ODE which is a polynomial in $\{ p, y \}$ must be a polynomial (with constant coefficients) of $\{ u_0^3(q^a), u_1^3(q^a) \}$.

Step 2. From the identity $(u_1^3)^2 = 2u_2^3 + 2qu_0^3$, one may choose as a basis for the vector space of such polynomial first integrals;

$$u_j^{m+1}(q^a) = (u_0^3)^{j_0} (u_1^3)^{j_1} (u_2^3)^{j_2}(q^a),$$

where

$$\begin{aligned} j_0 + j_1 + j_2 = m - 1 \geq 1, \\ j_0,\, j_2 \geq 0, \quad j_1 = 0, \text{ or } 1, \\ j = 0 \cdot j_0 + 1 \cdot j_1 + 2 \cdot j_2 \leq 2m-2. \end{aligned}$$

Note that each pair $(m,\, j)$, $m \geq 2$, $0 \leq j \leq 2m-2$, uniquely determines $(j_0,\, j_1,\, j_2)$.

Step 3. We claim that

$$\begin{aligned} \text{weight}\,(u_j^{m+1}) &= \ j_0 + j_1 + 2j_2, \\ \text{depth}\,(u_j^{m+1}) &= 2j_0 + j_1 + 2j_2. \end{aligned} \tag{4.39}$$

One checks that $(u_0^3,\, u_1^3,\, u_2^3)$ has weight $(1,\, 1,\, 2)$, and depth $(2,\, 1,\, 2)$ respectively, and the claim follows by induction.

Since $j_1 = 0$, or 1, the condition $\text{depth}\,(u_j^{m+1}) \leq (2d-4)+1$ translates to the condition $m \leq d$.

Step 4. Note that weight and depth uniquely determine $(m,\, j)$. It follows from Lemma 3.9 and the analysis in Section 3 that it suffices to show, for each $(m,\, j)$, $2 \leq m \leq d$, $0 \leq j \leq 2m-2$, that there are at most $d-m$ independent Abelian relations of the form

$$(\, v_1\, u_j^{m+1}(q^1),\ v_2\, u_j^{m+1}(q^2),\ \ldots\ ,\ v_d\, u_j^{m+1}(q^d)\,),$$

for constant coefficients $(v_1,\, v_2,\, \ldots v_d)$ (this implies $\dim \mathcal{S}_d = \rho_d$).

From Lemma 3.9 again, $u_j^{m+1}(q)$ is homogeneous of weight j ("weight" used in Section 3), and has degree $m-1$ in the indeterminate variable q. One may check that each of m coefficients of $u_j^{m+1}(q)$ as a polynomial in q is nonzero (see (3.6)), and hence linearly independent functions.[9] By Vandermonde identity (since $q^a \neq q^b$ for $a \neq b$), the coefficient $(v_1,\, v_2,\, \ldots v_d)$ as a vector in $\mathbb{R}^d$ must satisfy m independent linear equations.

Step 5. From Step 4, only those u_j^{m+1}, $2 \leq m \leq d-1$, give rise to a nonzero Abelian relation. By (4.39), $\text{depth}\,(u_j^{m+1}) = 2j_0 + j_1 + 2j_2 \leq 2j_0 + 2j_1 + 2j_2 = 2m-2 \leq 2d-4$. $\square$

Proof of Theorem 4.37. If the $(d-1)d$ linear constraint equations of depth $2d-3$ in the table above were degenerate, there must exist a nonzero Abelian relation of depth $2d-3 = (2d-4)+1$. $\square$

[9] Evaluate $u_0^3,\, u_1^3,\, u_2^3$ at $(x,\, y,\, p) = (-2, -1, 2)$, then each of them are of the form $\epsilon_0 + q\epsilon_1$, with $\epsilon_0,\, \epsilon_1 > 0$.

5. Legendrian 3-webs of maximum rank

In this section, we give a local analytic characterization of the Legendrian 3-webs of maximum rank three, and determine their local normal forms, Proposition 5.6, Theorem 5.9. We apply this result to the Legendrian 3-web associated with a pair of Riemannian metrics, Theorem 5.18.

In planar web geometry there exists a unique planar 3-web of maximum rank one up to diffeomorphism, which is locally equivalent to a set of three families of parallel lines in the plane.[10] In contrast, the results of this section show that the local moduli space of Legendrian 3-webs of maximum rank is generally two dimensional, and there already exist many non-trivial examples.

5.1. **Local normal forms.**

5.1.1. *Equivalence problem for Legendrian 3-webs.* In this sub-section, we employ the equivalence method of E. Cartan and determine the local structure equation for the Legendrian 3-webs in a contact three manifold.

Let X be a contact three manifold. Let $\mathcal{C} = \langle\, \theta \,\rangle$ be the contact line bundle generated by a contact 1-form θ. Let $\mathcal{I}^a \subset T^*X$, $a = 1, 2, 3$, be the rank 2 sub-bundles that define a Legendrian 3-web $\mathcal{W}$. Up to scaling and adding multiples of θ, one may arrange the generators $\mathcal{I}^a = \langle\, \theta;\, \theta^a \,\rangle$ such that

$$\theta^1 + \theta^2 + \theta^3 = 0. \tag{5.1}$$

Let B_0 be the sub-bundle of the principal frame bundle of X defined by the relation (5.1). We continue to use $\theta;\ \theta^a$, $a = 1, 2, 3$, to denote the corresponding tautological 1-forms on B_0. The structure group of the bundle B_0 acts on $\theta;\ \theta^a$, $a = 1, 2, 3$, by

$$(\,\theta,\, \theta^a\,) \to (\,\epsilon'\,\theta,\ \epsilon\,\theta^{\sigma(a)} + t^a\,\theta\,),$$

for nonzero scaling constants $\epsilon',\ \epsilon$, a permutation $\sigma \in S_3$, the symmetric group on three letters, and for translating constants $t^a,\ t^1 + t^2 + t^3 = 0$.

From the general theory, the tautological 1-forms satisfy the following structure equation.

$$d\begin{pmatrix}\theta^1\\ \theta^2\\ \theta\end{pmatrix} = -\begin{pmatrix}\alpha & \cdot & \beta^1\\ \cdot & \alpha & \beta^2\\ \cdot & \cdot & \beta\end{pmatrix}\wedge\begin{pmatrix}\theta^1\\ \theta^2\\ \theta\end{pmatrix} + \begin{pmatrix}s^1\,\theta^1\wedge\theta^2\\ s^2\,\theta^1\wedge\theta^2\\ s\,\theta^1\wedge\theta^2\end{pmatrix}. \tag{5.2}$$

Here $\alpha,\ \beta^1,\ \beta^2,\ \beta$ are pseudo-connection forms, and $s,\ s^1,\ s^2$ are torsion coefficients. It is easily checked that one may use the group action along the fibers of $B_0 \to X$ to normalize so that

$$s = 1,\ s^1 = s^2 = 0.$$

After modifying the pseudo-connection 1-form α as necessary, one consequently arrives at a sub-bundle $B \subset B_0$ on which the following structure equation holds.[11]

$$d\begin{pmatrix}\theta^1\\ \theta^2\\ \theta\end{pmatrix} = -\begin{pmatrix}\alpha & \cdot & \cdot\\ \cdot & \alpha & \cdot\\ \cdot & \cdot & 2\alpha\end{pmatrix}\wedge\begin{pmatrix}\theta^1\\ \theta^2\\ \theta\end{pmatrix} + \begin{pmatrix}\theta\wedge(R\,\theta^1 + S\,\theta^2)\\ \theta\wedge(T\,\theta^1 - R\,\theta^2)\\ \theta^1\wedge\theta^2\end{pmatrix}. \tag{5.4}$$

Here $R,\ S,\ T$ are new torsion coefficients. The exterior derivative identity $d(d(\theta)) = 0$ implies

$$d\alpha = \theta\wedge(\,N\theta^1 + L\,\theta^2\,), \tag{5.5}$$

for coefficients $N,\ L$.

[10] See [1][29] for the geometry associated with the hexagonal planar 3-webs.

[11] The structure group of the principal bundle B acts on $\theta;\ \theta^a$, $a = 1, 2, 3$, by $(\,\theta,\, \theta^a\,) \to (\,\mathrm{sgn}(\sigma)\epsilon^2\,\theta,\, \epsilon\,\theta^{\sigma(a)}\,)$, for a nonzero scaling constant ϵ, and a permutation by $\sigma \in S_3$. For example, under the permutation $(\,\theta^1,\, \theta^2,\, \theta^3) \to (\,\theta^2,\, \theta^1,\, \theta^3)$, the torsion coefficients transform to

$$\begin{pmatrix}R & S\\ T & -R\end{pmatrix} \to \begin{pmatrix}R & -T\\ -S & -R\end{pmatrix}. \tag{5.3}$$

Proposition 5.6. *Let $\mathcal{W}$ be a Legendrian 3-web on a contact three manifold X. There exists a sub-bundle B of the $\mathrm{GL}_3\mathbb{R}$ principal frame bundle of X on which the tautological 1-forms θ; θ^a, $a = 1, 2, 3$, satisfy the structure equations* (5.1), (5.4), (5.5). *The functional relations among the torsion coefficients R, S, T, N, L and their successive derivatives are the basic local invariants of $\mathcal{W}$. The algebra of local symmetry vector fields of $\mathcal{W}$ is at most four dimensional, and it is four dimensional whenever the torsion coefficients R, S, T, N, L all vanish.*

Proof. The algebra of local symmetry vector fields of $\mathcal{W}$ is four dimensional when R, S, T, N, L are all constants. Since they scale along the fibers of the bundle B, e.g., $dR \equiv 2R\alpha, \mod \theta, \theta^1, \theta^2$, this occurs only when R, S, T, N, L all vanish. The rest follows from the general theory of equivalence method. We omit the details. □

It will be shown in the next sub-section that the condition for a Legendrian 3-web to admit maximum number of linearly independent Abelian relations is expressed as a set of first order differential equations for the coefficients R, S, T, N, L.

Proposition 5.6 implies the following corollary on the intersection of the local algebra of symmetry vector fields of two distinct path geometries on a surface. Let M be a two dimensional manifold. Let $\mathbb{P}(TM) \to M$ be the projective tangent bundle equipped with the canonical contact structure. Recall that a path geometry on M is a (local) Legendrian foliation on $\mathbb{P}(TM)$ transversal to the fibers of the projection $\mathbb{P}(TM) \to M$, [7].

Corollary 5.7. *Let $\mathcal{F}_\pm$ be a pair of distinct path geometries on a two dimensional manifold M. Let $\mathfrak{p}_\pm \subset \mathfrak{X}(M)$ be the algebra of symmetry vector fields for $\mathcal{F}_\pm$ respectively (here $\mathfrak{X}(M)$ is the algebra of vector fields on M). Then*

$$\dim \mathfrak{p}_+ \cap \mathfrak{p}_- \leq 4.$$

The equality holds whenever the pair $\mathcal{F}_\pm$ is locally equivalent to the pair of path geometries defined by the second order ODE's

$$y'' = q_\pm, \qquad q_+,\, q_- \text{ are distinct constants.}$$

Proof. The prolongation of the intersection $\mathfrak{p}_+ \cap \mathfrak{p}_-$ to $\mathbb{P}(TM)$ is the symmetry algebra of vector fields for the Legendrian 3-web defined by $\mathcal{F}_\pm$, and the fibers of the projection $\mathbb{P}(TM) \to M$. □

5.1.2. *Abelian relations.* In this sub-section, the general computational algorithm for Abelian relations established in Section 4 is applied to the case of Legendrian 3-webs of maximum rank three. We give the local normal forms for such 3-webs in terms of a set of second order ODE's.

We continue the analysis of Section 5.1.

Let

$$\begin{aligned}\Omega^1 &= f\,\theta^1 + g^1\,\theta,\\ \Omega^2 &= f\,\theta^2 + g^2\,\theta,\\ \Omega^3 &= -\Omega^1 - \Omega^2.\end{aligned}$$

denote the sections of $\mathcal{I}^a$, $a = 1, 2, 3$, that satisfy the additive relation $\sum_b \Omega^b = 0$. The covariant derivatives of f, g^1, g^2 are denoted by

$$\begin{aligned}df &= f\,\alpha + f_1\theta^1 + f_2\theta^2 + f_0\theta,\\ dg^1 &= 2g^1\,\alpha + g^1_1\theta^1 + g^1_2\theta^2 + g^1_0\theta,\\ dg^2 &= 2g^2\,\alpha + g^2_1\theta^1 + g^2_2\theta^2 + g^2_0\theta.\end{aligned}$$

We shall adopt the similar notational convention for the covariant derivatives of the structure coefficients R, S, T, N, L, e.g., $dR = 2R\,\alpha + R_1\,\theta^1 + R_2\,\theta^2 + R_0\,\theta$, etc.

The defining equation $d\Omega^a = 0,\ a = 1,\ 2$, is equivalent to that

$$f_2 = g^1,\ f_1 = -g^2,$$
$$g^1_2 = Sf,\ g^2_1 = Tf,$$
$$g^1_1 = f_0 + Rf,\ g^2_2 = f_0 - Rf.$$

With this relation, the exterior derivative $d(d(f)) \wedge \theta = 0$ gives $f_0 = 0$. $d(d(f)) = 0$ now gives

$$g^1_0 = -Lf + Sg^2 + Rg^1,$$
$$g^2_0 = Nf - Rg^2 + Tg^1.$$

The linear differential system for the coefficients $\{ f,\ g^1,\ g^2 \}$ closes up at this step.

The exterior derivative identities $d(d(g^1)) = 0,\ d(d(g^2)) = 0$ give the following three compatibility equations for a Legendrian 3-web to admit an Abelian relation,

$$(S_0 + L_2)f + (-S_1 + 4L)g^1 - S_2 g^2 = 0,$$
$$(R_0 + L_1)f + (2N - R_1)g^1 + (-S_1 - L)g^2 = 0,$$
$$(-T_0 + N_1)f + T_1 g^1 + (-3N - R_1)g^2 = 0.$$

In order for the 3-web to have maximum rank three, these equations should hold identically and each of the nine coefficients of $f,\ g^1,\ g^2$ in the above must vanish, otherwise the rank is strictly less than three. This implies, after a short computation, that the structure coefficients of a Legendrian 3-web of maximum rank must satisfy

$$dR \equiv dS \equiv dT \equiv 0, \mod \alpha, \tag{5.8}$$
$$N = L = 0.$$

Take a section of the frame for which $\alpha = 0$. Then $R,\ S,\ T$ are constants, and (5.4) becomes the structure equation of a Lie group.[12] Hence *a Legendrian 3-web of maximum rank three is necessarily homogeneous*.

Theorem 5.9. *Let X be a contact three manifold. Let $\mathcal{W}$ be a Legendrian 3-web on X of maximum rank three. Then $\mathcal{W}$ is locally equivalent to a left invariant homogeneous Legendrian 3-web on a three dimensional Lie group. There exists a local contact isomorphism of X with the 1-jet space $J^1(\mathbb{R},\ \mathbb{R})$ equipped with the canonical contact structure such that $\mathcal{W}$ is defined by the integral curves of the following set of second order ODE's. Note that $R,\ S,\ T$'s in the expressions below are the (constant) structure coefficients of* (5.4) *with $\alpha = 0$.*

• ***Case*** $R^2 + ST = 0$***.*** *When $R = S = 0$,*

$$y'' + Ty = 0,$$
$$y'' + Ty + 1 = 0,$$

and the Legendrian foliation by the fibers of the projection $J^1(\mathbb{R},\ \mathbb{R}) \to J^0(\mathbb{R},\ \mathbb{R})$.

The general cases are obtained by applying the linear action (5.11) *to* (5.12)*.*

• ***Case*** $R^2 + ST > 0$***.*** *When $R > 0,\ S = T = 0$,*

$$y'' + R(y')^2 = 0, \tag{5.10}$$
$$y'' - R(y')^2 = 0,$$
$$y'' + R\tanh(Ry)(y')^2 = 0.$$

The general cases are obtained by applying the linear action (5.11) *to* (5.13)*.*

• ***Case*** $R^2 + ST < 0$***.***

[12] See [6, p29] for a classification of three dimensional Lie groups.

When $R=0,\ -S=T<0$*,*

$$\begin{aligned} y'' - T\tan(Ty)(y')^2 &= 0,\\ y'' + T\cot(Ty)(y')^2 &= 0,\\ y'' + T\frac{\cos(Ty)-\sin(Ty)}{\cos(Ty)+\sin(Ty)}(y')^2 &= 0. \end{aligned}$$

When $R=0,\ -S=T\neq 0$ *generally,*

$$\begin{aligned} y'' + \tan(Ty)(1-T(y')^2) &= 0,\\ y'' - \cot(Ty)(1-T(y')^2) &= 0,\\ y'' - \frac{\cos(Ty)-\sin(Ty)}{\cos(Ty)+\sin(Ty)}(1-T(y')^2) &= 0. \end{aligned}$$

The general cases are obtained by applying the linear action (5.11) *to* (5.14), (5.15).

Proof. The proof is by direct computation. Set the contact 1-form $\theta = dy - pdx$. Then θ and θ^1, θ^2 below satisfy the structure equation (5.4), (5.8) with $\alpha = 0$. Note that under the linear action by a matrix $\mathrm{g}\in \mathrm{SL}_2\mathbb{R}$,

$$\begin{pmatrix}\theta^1\\ \theta^2\end{pmatrix} \to \mathrm{g}\begin{pmatrix}\theta^1\\ \theta^2\end{pmatrix}, \quad \text{then} \quad \begin{pmatrix}R & S\\ T & -R\end{pmatrix} \to \mathrm{g}\begin{pmatrix}R & S\\ T & -R\end{pmatrix}\mathrm{g}^{-1}. \tag{5.11}$$

• **Case** $R^2+ST=0$**.** Up to the conjugation (5.11), one may assume either $R=S=0, T\neq 0$, or $R=S=T=0$. Set

$$\begin{aligned} \theta^1 &= dx,\\ \theta^2 &= dp + Tydx,\\ \theta^1+\theta^2 &= dp + (Ty+1)dx. \end{aligned} \tag{5.12}$$

• **Case** $R^2+ST>0$**.** Up to the conjugation (5.11), one may assume $R>0,\ S=T=0$. Set

$$\begin{aligned} \theta^1 &= \frac{\exp(Ry)}{\sqrt{2R}\,p}(dp + Rp^2dx),\\ \theta^2 &= \frac{\exp(-Ry)}{\sqrt{2R}\,p}(dp - Rp^2dx),\\ \theta^1+\theta^2 &= \frac{2\cosh(Ry)}{\sqrt{2R}p}(dp + R\tanh(Ry)p^2dx). \end{aligned} \tag{5.13}$$

• **Case** $R^2+ST<0$**.** Up to the conjugation (5.11), one may assume either $R=0,\ -S=T<0$, or $R=0,\ -S=T>0$.

In case $T<0$, set

$$\begin{aligned} \theta^1 &= \frac{\cos(Ty)}{\sqrt{-T}p}(dp - T\tan(Ty)p^2dx),\\ \theta^2 &= \frac{\sin(Ty)}{\sqrt{-T}p}(dp + T\cot(Ty)p^2dx),\\ \theta^1+\theta^2 &= \frac{\cos(Ty)+\sin(Ty)}{\sqrt{-T}p}(dp + T\frac{\cos(Ty)-\sin(Ty)}{\cos(Ty)+\sin(Ty)}p^2\,dx). \end{aligned} \tag{5.14}$$

In the general case, set (this also works for the case $T<0$)

$$\theta^1 = -\frac{\cos(Ty)}{\sqrt{1-Tp^2}}(dp+\tan(Ty)(1-Tp^2)dx), \tag{5.15}$$
$$\theta^2 = -\frac{\sin(Ty)}{\sqrt{1-Tp^2}}(dp-\cot(Ty)(1-Tp^2)dx),$$
$$\theta^1+\theta^2 = -\frac{\cos(Ty)+\sin(Ty)}{\sqrt{1-Tp^2}}(dp-\frac{\cos(Ty)-\sin(Ty)}{\cos(Ty)+\sin(Ty)}(1-Tp^2)dx). \square$$

Consider the case $R=1$, $S=T=0$. The general solutions to the given set of ODE's (5.10) are

$$y = \ \ \ln(\,\epsilon_1^1 x+\epsilon_2^1\,),$$
$$y = -\ln(\,\epsilon_1^2 x+\epsilon_2^2\,),$$
$$y = \text{arcsinh}(\,\epsilon_1^3 x+\epsilon_2^3\,), \quad \epsilon_j^i\text{'s are constants.}$$

The three Abelian relations are

$$\big(\ e^y y', \qquad e^{-y}y', \qquad -2\cosh(y)y'\ \big),$$
$$\big(\ e^y(xy'-1), \qquad e^{-y}(xy'+1), \qquad -2\cosh(y)xy'+2\sinh(y)\ \big),$$
$$\big(\ e^y(x^2y'-2x+\frac{1}{y'}), \quad e^{-y}(x^2y'+2x+\frac{1}{y'}), \quad -2\cosh(y)x^2y'+4x\sinh(y)-\frac{2\cosh(y)}{y'}\big).$$

The Abelian relations for the other cases can be determined in a similar way by direct computation.

5.2. **A pair of geodesic foliations with Abelian relations.** In this section, we give an application of Proposition 5.6 to the case the Legendrian 3-web is defined by the geodesic foliations of a pair of two dimensional Riemannian metrics, Definition 5.16. Proposition 5.6 implies that such a Legendrian 3-web has maximum rank three whenever the pair of metrics share three dimensional algebra of projective vector fields. The classification result of [4] then immediately gives the local normal forms for such pair of metrics, Theorem 5.9.

Let M be a two dimensional manifold equipped with a Riemannian metric g. Let $\mathbb{P}(TM)\to M$ be the projective tangent bundle with the canonical contact structure. The geodesic foliation of g on $\mathbb{P}(TM)$ is by definition a Legendrian foliation. A finite set of Riemannian metrics on M in this way give rise to a geodesic Legendrian web on $\mathbb{P}(TM)$.

Definition 5.16. *Let $g_\pm$ be a pair of Riemannian metrics on a two dimensional manifold M. Under an appropriate transversality condition,*[13] the Legendrian 3-web $\mathcal{W}_{g_\pm}$ on $\mathbb{P}(TM)$ associated with $g_\pm$ *is the 3-web defined by the geodesic foliations of $g_\pm$, and the fibers of projection $\mathbb{P}(TM)\to M$. The pair of Riemannian metrics are* maximally geodesically compatible *when the associated Legendrian 3-web $\mathcal{W}_{g_\pm}$ has maximum rank three.*

Let us recall some of the relevant facts on the two dimensional Riemannian metrics. We refer to [4][24][23] for the details.

Let M be a two dimensional oriented Riemannian manifold with the given metric g. Let $\pi: FM\to M$ be the SO_2-bundle of oriented orthonormal frames. Let $\{\,\omega^1,\,\omega^2\,\}$ be the canonical 1-forms on FM such that $\pi^*g=(\omega^1)^2+(\omega^2)^2$. The Levi-Civita connection 1-form ρ, and the Gauss curvature function K are uniquely determined by the structure equation

$$d\omega^1 = -\rho\wedge\omega^2, \tag{5.17}$$
$$d\omega^2 = \ \ \rho\wedge\omega^1,$$
$$d\rho = K\,\omega^1\wedge\omega^2.$$

[13] See **Step 1** in Section 5.2.1.

Differentiating the last equation, the higher order covariant derivatives of K are inductively defined by

$$\begin{aligned} dK &= K_1\omega^1 + K_2\omega^2, \\ dK_1 &= -K_2\rho + K_{11}\omega^1 + K_{12}\omega^2, \\ dK_2 &= \quad K_1\rho + K_{21}\omega^1 + K_{22}\omega^2, \quad K_{12} = K_{21}, \quad \text{etc.} \end{aligned}$$

The set of un-parameterized geodesics of g defines the geodesic flow (foliation) $\mathcal{F}_g$ on $\mathbb{P}(TM)$, which is naturally extended to the tangent bundle $TM \to M$. Let $S^k(T^*M) \to M$ be the bundle of symmetric k-forms on M considered as functions on TM which are homogeneous of degree k on the fibers. Set

$$\begin{aligned} I_k(g) &= \{\, \sigma \in C^\infty(S^k(T^*M)) \mid \ \sigma \ \text{ is a first integral for } \mathcal{F}_g \ \}, \\ \varrho_k(g) &= \text{dimension}_{\mathbb{R}}\, I_k(g). \end{aligned}$$

The dimensions $\varrho_k(g)$, $k = 1, 2, \ldots$, are the basic integral invariants of a two dimensional Riemannian metric g.

The following results on $I_k(g)$ for $k = 1, 2, 3$, are known. The class of Riemannian metrics that are of interest to us are the so called *Darboux super-integrable* metrics, for which $\varrho_2(g) = 4$.

$I_1(g)$:
An element of $I_1(g)$ corresponds to a Killing vector field of g. For any metric g, $\varrho_1(g) = 3$, 1, or 0 (generic case);

$\varrho_1(g) = 3$ when g is a metric of constant curvature.

$\varrho_1(g) = 1$ when g admits a Killing vector field, which is unique up to scale. An analytic characterization of this class of metrics is given in [15]. For example, the curvature must satisfy the second order equation $K_1K_2(K_{11} - K_{22}) - (K_1^2 - K_2^2)K_{21} = 0$.

$I_2(g)$:
An element of $I_2(g)$ corresponds to a Riemannian metric projectively equivalent to g, [23]. For any metric g, $\varrho_2(g) = 6, 4, \ldots$, or 1(generic case);

$\varrho_2(g) = 6$ when g is a metric of constant curvature.

$\varrho_2(g) = 4$ when g admits a Killing field ($\varrho_1(g) = 1$), and the curvature of g satisfies an additional set of second, and third order differential equations. This class of metrics are called *Darboux super-integrable*, [4].

Darboux super-integrable metrics are equivalently characterized as having three dimensional algebra of projective vector fields. [4] gave, among other things, the explicit local normal forms for Darboux super-integrable metrics combining this observation with a classical result of Lie on the classification of the algebra of projective vector fields on the plane.

$I_3(g)$:
The case of cubic integrals are studied in [24]. For any metric g, $\varrho_3(g) \leq 10$, and $\varrho_3(g) = 10$ when g is a metric of constant curvature. It is possible that the next admissible value of $\varrho_3(g)$ is 4, which is attained by the Darboux super-integrable metrics.

The main result of this section is that a two dimensional Riemannian metric g_+ admits a maximally geodesically compatible mate g_- when g_+ is either of constant curvature, or Darboux super-integrable. Note from the remark above Theorem 5.9 that a Riemannian metric admitting a maximally geodesically compatible mate has at least three dimensional algebra of projective vector fields.

Theorem 5.18. *Let g_+ be a two dimensional Riemannian metric. g_+ admits a maximally geodesically compatible mate g_- whenever g_+ is either of constant curvature, or Darboux super-integrable. Let $\mathcal{R}_{g_+}$ denote the moduli space of Riemannian metrics maximally geodesically compatible with the given metric g_+. Then,*

a) when g_+ is of constant curvature, $\mathcal{R}_{g_+}$ consists of 10 parameter family of metrics of constant curvature, and 8 parameter family of Darboux super-integrable metrics.

b) when g_+ is Darboux super-integrable, $\mathcal{R}_{g_+}$ consists of 5 parameter family of metrics of constant curvature, and 4 parameter family of Darboux super-integrable metrics.

Let us mention that many of the metrics in this theorem are geodesically equivalent. For example, for the Darboux super-integrable metric g_+ considered in Section 5.2.2, all of the 5 parameter family of mates of constant curvature are geodesically equivalent, and the 4 parameter family of Darboux super-integrable mates are *foliated* by 3 parameter families of geodesically equivalent metrics.

5.2.1. *Differential analysis.* The differential analysis for the proof of Theorem 5.18 is a straightforward application of the over-determined PDE machinery. Due to the complexity and size of the algebraic expressions involved, the computations were performed on the computer algebra system `Maple`. Let us record the relevant steps of the analysis, only for the case of Darboux super-integrable metrics. The analysis for the constant curvature metrics is similar, and shall be omitted.

Given a Darboux super-integrable metric g_+ on a two dimensional manifold M, let $FM \to M$ be the associated oriented orthonormal frame bundle. Let $\{\, \omega^1,\, \omega^2,\, \rho \,\}$ be the structure forms on FM that satisfy the structure equation (5.17). From the general theory, the geodesic foliation of g_+ on FM is defined by the rank 2 sub-bundle

$$\mathcal{I}^+ = \langle\, \omega^2,\, \rho \,\rangle \subset T^*FM.$$

Here ω^2 defines the canonical contact structure on FM.

Step 0. Let g_- be another metric on M. When pulled back to FM, one may write

$$g_- = (\eta^1)^2 + (\eta^2)^2,$$

for a g_--orthonormal coframe $\{\, \eta^1,\, \eta^2 \,\}$, where

$$\begin{aligned} \eta^1 &= a\,\omega^1 + b\,\omega^2, \\ \eta^2 &= \qquad\quad c\,\omega^2, \end{aligned} \tag{5.19}$$

for variables $a,\, b,\, c$. Let ψ denote the connection 1-form for $\{\, \eta^1,\, \eta^2 \,\}$ so that

$$\begin{aligned} d\eta^1 &= -\psi \wedge \eta^2, \\ d\eta^2 &= \quad \psi \wedge \eta^1, \\ d\psi &= Q\,\eta^1 \wedge \eta^2, \end{aligned} \tag{5.20}$$

where Q is the curvature of the metric g_-.

Differentiating (5.19), the structure equation (5.20) implies that

$$\begin{aligned} da &= -b\rho \qquad + a_1\omega^1 + a_2\omega^2, \\ db &= a\rho - c\psi + b_1\omega^1 + b_2\omega^2, \\ dc &= \quad b\psi \qquad + c_1\omega^1 + c_2\omega^2, \end{aligned}$$

where $a_2 = b_1$. Here $a_j,\, b_j,\, c_j$'s denote the covariant derivatives of $a,\, b,\, c$. The connection form ψ is given by

$$\psi = \frac{1}{a}(c\rho + t\omega^1 - c_1\omega^2),$$

for a variable t. Differentiating this equation, one may write

$$dt = 2c_1\rho - a_1\psi + t_1\omega^1 + t_2\omega^2.$$

Let us remark here that we shall adopt the similar notational convention for the rest of the analysis, i.e., $da_1 \equiv a_{11}\omega^1 + a_{12}\omega^2, \mod \rho,\, \psi$, etc. The curvature Q for the metric g_- is for example given by

$$Q = \frac{-c_{11} - t_2 + Kc}{a^2 c}. \tag{5.21}$$

Step 1. Set the rank 2 sub-bundles

$$\begin{aligned}\mathcal{I}^0 &= \langle\, \omega^2,\, \omega^1 \,\rangle,\\ \mathcal{I}^- &= \langle\, \eta^2,\, \psi \,\rangle,\\ &= \langle\, \omega^2,\, c\rho + t\omega^1 \,\rangle.\end{aligned}$$

Then the set of three foliations defined by $\{\,\mathcal{I}^{\pm},\, \mathcal{I}^0\,\}$ is the Legendrian 3-web $\mathcal{W}_{g_\pm}$ on FM, under the non-degeneracy assumption that $a,\, c,\, t \neq 0$, which we assume from now on.

Step 2. Following the analysis of Section 5.1, it is a straightforward computation to determine the associated coframe $\theta,\, \theta^a,\, a = 1,\, 2,\, 3$, and α, for $\mathcal{W}_{g_\pm}$ that satisfy the structure equation (5.4), (5.5). The torsion coefficients $R,\, S,\, T,\, N,\, L$ are then determined as rational functions in the successive derivatives of $a,\, b,\, c,\, t$.

Step 3. The claim is that the integrability condition (5.8) for the Legendrian 3-web $\mathcal{W}_{g_\pm}$ to have maximum rank three allows one to close up the structure equation for $a,\, b,\, c,\, t$.

Differentiating R and evaluating modulo $\alpha,\, \omega^2$, one may solve for $\{\,c_{21},\, t_{21}\,\}$ in terms of the rest of the variables. Differentiating S and evaluating modulo $\alpha,\, \omega^2,\, \omega^1$ with these relations, one may solve for $\{\,b_2\,\}$. Successively differentiating this equation for b_2, one may solve for $\{\,b_{21},\, b_{22},\, c_2\,\}$. Successively differentiating again the equation for c_2, one may solve for $\{\,b_{11},\, c_{22}\,\}$.

Differentiating R and evaluating modulo $\alpha,\, \omega^1,\, \rho$, one may solve for $\{\,t_{22}\,\}$. This implies that $dR,\, dS \equiv 0,\ \mathrm{mod}\ \alpha$.

Step 4. At this step, the remaining undetermined second derivatives of $a,\, b,\, c,\, t$ are $a_{11},\, c_{11},\, t_{11}$. Introduce the third order derivatives $da_{11} \equiv a_{111}\omega^1 + a_{112}\omega^2,\ \mathrm{mod}\ \rho$, etc, for these variables.

From the exterior derivative identities $d(d(a_1)),\, d(d(c_1)),\, d(d(t_1)) = 0$, one may solve for $\{\,a_{112},\, c_{112},\, t_{112}\,\}$. From the exterior derivative identity $d(d(b_1)) = 0$, one may solve for $\{\,c_{111}\,\}$. Differentiating T and evaluating modulo $\alpha,\, \omega^2$ with these relations, one may solve for $\{\,a_{111},\, t_{111}\,\}$. The structure equations for $a,\, b,\, c,\, t$ now close up at this step. We remark that one already has $d\alpha = 0$.

Step 5. A key integrability equation is obtained from the identity $d(d(\frac{c}{a}a_{11} - 2c_{11})) = 0$, which gives

$$3\,atK_1 + (tb + ca_1 - 2\,ac_1)K_2 = 0.$$

Assuming the metric g_+ is not of constant curvature, one may solve for $\{\,c_1\,\}$. Differentiating this equation for c_1, one may solve for $\{\,c_{11}\,\}$.

Step 6. Differentiating T and evaluating modulo α, one may solve for $\{\,t_{11}\,\}$. Differentiating this equation for t_{11}, one may solve for $\{\,t_1,\, t_2\,\}$. Note that the set of remaining independent variables at this step are $\{\,a,\, b,\, c,\, t;\, a_1,\, b_1,\, a_{11}\,\}$.

Taking the exterior derivative $d(d(a_{11})) = 0$, the resulting single integrability equation factors into two parts. The vanishing of the one part is equivalent to that the curvature Q of the metric g_- is constant. Assuming this is not the case, the vanishing of the other integrability equation allows one to solve for $\{\,b_1\,\}$. Differentiating this equation for b_1, one may solve for $\{\,a_{11}\,\}$.

At this stage, the remaining independent variables are $\{\,a,\, b,\, c,\, t;\, a_1\,\}$, and the structure equations for these variables are compatible. Since the metric g_- is well defined on the surface M, and is invariant under the SO_2 action along the fibers of $FM \to M$, from the general theory it follows that the moduli space $\mathcal{R}_{g_+}$ is generically $5 - 1 = 4$ dimensional.

The analysis for the case when the metric g_- has constant curvature follows the similar path. □

5.2.2. *Example.* Lie classified the possible local symmetry Lie algebras, and the representations thereof, of projective vector fields on the plane. Bryant, Manno, & Matveev used this classification to give explicit local normal forms for the two dimensional (pseudo) Riemannian metrics admitting a transitive algebra of projective vector fields, [4]. In this sub-section, we apply these results to our setting and give explicit examples of the maximally geodesically compatible pairs of Darboux super-integrable metrics. We shall closely follow the notations in [4].

Let $(\,x,\, y\,)$ be a local coordinate of $\mathbb{R}^2$. Let g_+ be the Darboux super-integrable metric

$$g_+ = e^{3x}dx^2 - 2D_+e^x dy^2. \tag{5.22}$$

The projective connection associated with g_+, or equivalently the equation of un-parameterized geodesics of g_+, is given by the second order ODE

$$y'' = \frac{1}{2}y' + D_+ e^{-2x}(y')^3. \tag{5.23}$$

The local symmetry algebra of (5.23) is generated by

$$\langle\ \frac{\partial}{\partial y},\ \frac{\partial}{\partial x} + y\frac{\partial}{\partial y},\ 2y\frac{\partial}{\partial x} + y^2\frac{\partial}{\partial y}\ \rangle. \tag{5.24}$$

One may verify by direct computation that the projective connection invariant under (5.24) is defined by the second order ODE's of the following form

$$y'' = \frac{1}{2}y' + De^{-2x}(y')^3, \quad D \text{ is constant}. \tag{5.25}$$

This projective connection is flat whenever $D = 0$.[14] Since we are considering the pair of Darboux super-integrable metrics, assume $D \neq 0$.

The maximally geodesically compatible mates of g_+ share the local symmetry algebra of projective vector fields. It thus suffices to find the metrics which has (5.25) as the equation of un-parameterized geodesics. Consider for example the metrics of the form

$$g_- = E(x,y)dx^2 + G(x,y)dy^2.$$

Then g_- is a mate of g_+ with the equation of geodesics (5.25) whenever

$$E(x,y) = E(x) = \frac{e^{(x+c_1)}}{(-2De^{(-x+c_1)} + c_2)^2},$$
$$G(x,y) = G(x) = \frac{1}{-2De^{(-x+c_1)} + c_2}, \quad D \neq D_+;\ c_1,\ c_2 \text{ are constants}.$$

The three (un-differentiated) Abelian relations for the Legendrian 3-web $\mathcal{W}_{g_\pm}$ are;

$$(\frac{-2D_+e^{-x}y^2p^2 + 4p^2e^x + e^xy^2 - 4pe^xy}{2p^2(D-D_+)},\ \frac{2De^{-x}y^2p^2 - 4p^2e^x - e^xy^2 + 4pe^xy}{2p^2(D-D_+)},\ -e^{-x}y^2),$$
$$(\frac{-2D_+e^{-x}yp^2 - 2pe^x + e^xy}{2p^2(D-D_+)},\ \frac{2De^{-x}yp^2 + 2pe^x - e^xy}{2p^2(D-D_+)},\ -e^{-x}y\),$$
$$(\frac{-2D_+e^{-x}p^2 + e^x}{2p^2(D-D_+)},\ \frac{2De^{-x}p^2 - e^x}{2p^2(D-D_+)},\ -e^{-x}\).$$

6. Concluding remarks

1. As stated in Section 1, the purpose of the present paper is to propose the Legendrian web for a second order generalization of the planar web. The list of dualities associated with the rank 2 simple Lie groups in Fig. 6.1 provides an algebro geometric perspective on this generalization, [8]. Here Z', Z, Z'' are the respective incidence spaces.

It is a natural extension to consider the analogous problem for the webs on $\mathbb{Q}^5$ defined by the horizontal foliations with respect to the $\mathrm{G}_2^{\mathbb{C}}$ invariant rank 2 distribution (Cartan distribution).

2. Given a set of $n+2$ points $\{\mathrm{p}^1, \mathrm{p}^2, \ldots \mathrm{p}^{n+2}\}$ in $\mathbb{RP}^n$ in general position, the associated *exceptional* $n+3$ *web* $\mathcal{E}_{n+3}$ consists of the $n+2$ bundles of lines with vertices p^a's, and the family of rational normal curves through p^a's. Damiano in his thesis gave a characterization that, up to diffeomorphism, $\mathcal{E}_{n+3}$ is the unique non-linearizable quadrilateral web of curves in dimension n, [14]. He also showed that $\mathcal{E}_{n+3}$ is of maximum rank based on the observation that $\mathcal{E}_{n+3}$ naturally occurs on the (smooth part of) configuration space of ordered set of

[14] This shows in particular that the five parameter family of metrics of constant curvature which are maximally geodesically compatible with the given metric g_+ are all geodesically equivalent.




Figure 6.1. Dualities associated with the rank 2 simple Lie groups.

$n+3$ points in $\mathbb{RP}^1$, which is essentially the quotient space of the Grassmannian $Gr(2, \mathbb{R}^{n+3})$ by the Cartan subgroup of $\mathrm{SL}_{n+3}\mathbb{R}$. Modulo the codimension one subspace of combinatorial Abelian relations, the remaining single non-combinatorial Abelian relation is given by the trace of the appropriate power of the harmonic form representing the Euler class of the canonical bundle of $Gr(2, \mathbb{R}^{n+3})$.

It would be interesting if such a geometric construction exists for Legendrian webs.

3. We currently do not have any nontrivial examples of Legendrian 4-webs of maximum rank eleven. Let us give a description of one candidate.

Let (x^1, x^2, x^3, x^4) be the coordinate of $\mathbb{R}^4$. Let $\varpi = dx^1 \wedge dx^2 + dx^3 \wedge dx^4$ be the standard symplectic 2-form. Let $\mathrm{S}^3 \subset \mathbb{R}^4$ be the unit sphere equipped with the U_2-invariant induced contact structure defined by the contact 1-form $\theta = -x^2dx^1 + x^1dx^2 - x^4dx^3 + x^3dx^4$. Consider the Legendrian 4-web $\mathcal{W}$ on a generic small open subset of S^3 defined by the rank 2 sub-bundles

$$\mathcal{I}^a = \{\, \theta,\ dx^a \,\}, \quad a = 1,\, 2,\, 3,\, 4.$$

Each of the four sub 3-webs of $\mathcal{W}$ has rank 0, and $\mathcal{W}$ is distinct from the examples in Section 3. Is $\mathcal{W}$ algebraic in any way?

APPENDIX

We record the `Maple` code to verify that the set of matrices in Theorem 4.24 are strongly nondegenerate.

Step 1. This procedure defines the coefficient $\mathtt{c}^{\mathtt{i}}_{\mathtt{j}}$.

```
> c:=proc(i,j)
    if j < 0 then 0
    elif j > i/2 then 0
    else i!/(2^j*((i-2*j)!)*j!)  fi:
  end:
```

Step 2. M[d,k,s] is defined as the upper-left corner s-by-s sub-matrix of the k-th matrix (of size d-k) from Theorem 4.24 whose (1,1)-entry is $\mathtt{c}^{\mathtt{d+k-1}}_{\mathtt{k}}$.

```
> for d from 3 to 100 do
      for k from 1 to d-1 do
         for s from 1 to d-k do
            M[d,k,s]:= Matrix(s, s, (i, j) -> c(d+k-i,k-i+j)):
         end do:
      end do:
  end do:
```

Step 3. This procedure checks the determinant of M[d,k,s].

```
> for d from d0 to d1 do
      for k from 1 to d-1 do
         for s from 1 to d-k do
            if det(M[d,k,s])=0
               then print([d,k,s])
```

```
            elif [d,k,s] = [d1, d1-1,1]
               then print([d,k,s])
            else fi:
         end do:
      end do:
end do:
```

The `Maple` computation confirms the claim for `d` up to 100.

REFERENCES

[1] Agafonov, S.I., *Flat 3-webs via semi-simple Frobenius 3-folds*, http://arxiv.org/abs/1108.1997 (2011), preprint

[2] Blaschke, W.; Bol, G., Geometrie der Gewebe, Topologische Fragen der Differentialgeometrie, Die Grundlehren der mathematischen Wissenschaften, Bd. XLIX, 1938

[3] Bryant, R.L.; Chern, S.-S.; Gardner, R.B.; Goldschmidt, H.L.; & Griffiths, Phillip, Exterior differential systems. MSRI Publications, 18. Springer-Verlag, 1991

[4] Bryant, Robert L.; Manno, Gianni; Matveev, Vladimir S., *A solution of a problem of Sophus Lie: normal forms of two-dimensional metrics admitting two projective vector fields*, Math. Ann. 340, No. 2 (2008), 437–463

[5] Bol, G., *On n-webs of curves in a plane*, Bull. Amer. Math. Soc. Volume 38, no 12 (1932), 855–857

[6] Bryant, R. L., *An introduction to Lie groups and symplectic geometry*, Geometry and quantum field theory. AMS IAS/Park City Math. Ser. 1 (1995), 5–181

[7] Bryant, R. L., *Projectively flat Finsler 2-spheres of constant curvature*, Selecta Math. 3, No.2 (1997), 161–203

[8] Bryant, R. L., *Élie Cartan and geometric duality*, Journées Élie Cartan 1998 et 1999, vol. 16 (2000), 5–20

[9] Cartan, E., *Sur les varietes a connexion projective*, Bull. Soc. Math. France 52 (1924), 205–241

[10] Chern, S.-S.; Griffiths, Phillip, *Abel's theorem and webs*, Jahr. Deutsch. Math. - Ver. 80 (1978), 13–110

[11] Chern, S.-S.; Griffiths, Phillip, *An inequality for the rank of a web and webs of maximum rank*, Ann. Sc. Norm. Super. Pisa, Cl. Sci., IV. Ser. 5 (1978), 539–557

[12] Chern, S.-S., *Abzählungen für Gewebe*, Abh. Math. Sem. Univ. Hamburg 11 (1935), 163–170

[13] Chern, S.-S., *Web geometry*, Bull. Am. Math. Soc., New Ser. 6 (1982), 1–8

[14] Damiano, D. B., *Webs, Abelian Equations and Characteristic Classes*, Thesis, Brown University (1980)

[15] Robert L. Foote, Chong-Kyu Han, Jong-Won Oh, *Infinitesimal isometries along curves and generalized Jacobi equations*, J. Geom. Anal., to appear

[16] Griffiths, P. A.; Harris, J., Principles of algebraic geometry, Pure and Applied Mathematics. Wiley-Interscience, New York, 1978

[17] Grifone, Joseph; Salem, Eliane (ed.), Web theory and related topics, World Scientific, Singapoer, 2001

[18] Gelfand, Israel M.; Zakharevich, Ilya, *Webs, Veronese curves, and bihamiltonian systems*, J. Funct. Anal. 99, No.1 (1991), 150–178

[19] Hénaut, Alain, *Formes différentielles abéliennes, bornes de Castelnuovo et géométrie des tissus*, Comment. Math. Helv. 79, no. 1 (2004), 25–57

[20] Henkin, Gennadi; Passare, Mikael, *Abelian differentials on singular varieties and variations on a theorem of Lie-Griffiths*, Invent. Math. 135, no. 2 (1999), 297–328

[21] Ivey, Thomas A.; Landsberg, J. M., Cartan for beginners, Graduate Studies in Mathematics, 61. AMS, 2003

[22] Little, John B., *On webs of maximum rank*, Geom. Dedicata, 31, no. 1 (1989), 19–35

[23] Matveev, Vladimir S., *Proof of the projective Lichnerowicz-Obata conjecture*, J. Differ. Geom. 75, No. 3 (2007), 459–502

[24] Matveev, Vladimir S.; Shevchishin, Vsevolod V., *Two-dimensional superintegrable metrics with one linear and one cubic integral*, J. Geom. Phys. 61, No. 8 (2011), 1353–1377

[25] Olver, Peter J., Equivalence, invariants, and symmetry, Cambridge: Cambridge University Press (1995)

[26] Pirio, Luc, *Abelian functional equations, planar web geometry and polylogarithms*, Selecta Math., New Ser. 11, No. 3-4 (2005), 453–489

[27] Pereira, J. V.; Pirio, Luc, An invitation to web geomety - From Abel's addition theorem to the algebraization of codimension one webs, IMPA Mathematical Publications, IMPA, Rio de Janeiro, 2009

[28] Trépreau, Jean-Marie, *Algebrisation des Tissus de Codimension 1 - La generalisation d'un Theoreme de Bol*, Inspired by Chern, Nankai Tracts in Mathematics 11 (2006), 399–433

[29] J. S. Wang , *On the Gronwall conjecture*, J. Geom. Anal., DOI: 10.1007/s12220-010-9184-6 (2010), online first

[30] J. S. Wang , *Projectively deformable Legendrian surfaces*, http://arxiv.org/abs/1107.4158 (2011), preprint

ST. LOUIS, MO 63124
E-mail address: `jswang12@gmail.com`